

# Lighting the way: Compelling open questions in photosynthesis research

Nancy A. Eckardt,<sup>1,\*</sup> Yagut Allahverdiyeva,<sup>2</sup> Clarisa E. Alvarez,<sup>3</sup> Claudia Büchel,<sup>4</sup> Adrien Burlacot,<sup>5,6</sup> Tanai Cardona,<sup>7,8</sup> Emma Chaloner,<sup>7,8</sup> Benjamin D. Engel,<sup>9</sup> Arthur R. Grossman,<sup>5,6</sup> Dvir Harris,<sup>10</sup> Nicolas Herrmann,<sup>4</sup> Michael Hodges,<sup>11</sup> Jan Kern,<sup>12</sup> Tom Dongmin Kim,<sup>7,8</sup> Veronica G. Maurino,<sup>13</sup> Conrad W. Mullineaux,<sup>7</sup> Henna Mustila,<sup>2</sup> Lauri Nikkanen,<sup>2</sup> Gabriela Schlau-Cohen,<sup>10</sup> Marcos A. Tronconi,<sup>3</sup> Wojciech Wietrzynski,<sup>9</sup> Vittal K. Yachandra,<sup>12</sup> Junko Yano<sup>12</sup>

<sup>1</sup>The Plant Cell, American Society of Plant Biologists, USA

<sup>2</sup>Molecular Plant Biology Unit, Department of Life Technologies, University of Turku, 20014 Turku, Finland

<sup>3</sup>Centro de Estudios Fotosintéticos y Bioquímicos (CEFOTI-CONICET), Facultad de Ciencias Bioquímicas y Farmacuticas, University of Rosario, Suipacha 570, 2000 Rosario, Argentina

<sup>4</sup>Institute of Molecular Biosciences, Goethe University Frankfurt, 60438 Frankfurt, Germany

<sup>5</sup>Division of Bioscience and Engineering, Carnegie Institution for Science, 260 Panama Street, Stanford, CA 94305, USA

<sup>6</sup>Department of Biology, Stanford University, Stanford, CA 94305, USA

<sup>7</sup>School of Biological and Behavioural Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, UK

<sup>8</sup>Department of Life Sciences, Imperial College London, London SW7 2AZ, UK

<sup>9</sup>Biozentrum, University of Basel, Sptialstrasse 41, 4056 Basel, Switzerland

<sup>10</sup>Department of Chemistry, Massachusetts Institute of Technology, Massachusetts Ave, Cambridge, MA 02139, USA

<sup>11</sup>Université Paris-Saclay, CNRS, INRAE, Université d'Evry, Université de Paris Cité, Institute of Plant Sciences Paris-Saclay (IPSP), 91190 Gif-sur-Yvette, France

<sup>12</sup>Molecular Biophysics and Integrated Biomaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>13</sup>Molecular Plant Physiology, Institute for Cellular and Molecular Botany (IZMB), University of Bonn, Kirschallee 1, 53115 Bonn, Germany

\*Author for correspondence: [neckardt@aspb.org](mailto:neckardt@aspb.org).

†Authors are listed alphabetically (except for the lead author/coordinating editor).

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plcell>) is: Nancy A. Eckardt ([neckardt@aspb.org](mailto:neckardt@aspb.org)).

## Abstract

Photosynthesis—the conversion of energy from sunlight into chemical energy—is essential for life on Earth. Yet there is much we do not understand about photosynthetic energy conversion on a fundamental level: how it evolved and the extent of its diversity, its dynamics, and all the components and connections involved in its regulation. In this commentary, researchers working on fundamental aspects of photosynthesis including the light-dependent reactions, photorespiration, and C<sub>4</sub> photosynthetic metabolism pose and discuss what they view as the most compelling open questions in their areas of research.

## Introduction

### (Written by Nancy A. Eckardt)

In the garden on a September morning, I was, once again, astonished to find that a tiny seed I planted a few months ago had produced large stems, leaves, and—seemingly daily—dozens of enormous fruit (also known as zucchini), literally out of thin air. This plant took carbon as CO<sub>2</sub> directly out of the air and turned it into organic matter, and along the way, split water molecules to produce oxygen and stored energy, in one of the most remarkable and essential life processes: photosynthesis.

All oxygenic photosynthetic organisms have two photosystems (PS) known as PSI and PSII, which are embedded in thylakoid membranes and carry out light-driven electron transport. PSII absorbs light energy of wavelengths <680 nm to oxidize water into molecular oxygen (the “water-splitting” reaction), which contributes to the proton gradient needed to convert ADP to ATP, whereas PSI absorbs energy of longer wavelengths to generate high-energy electrons used to convert NADP<sup>+</sup> to NADPH. All PSs have an array of light-harvesting antennae and mechanisms to dissipate excess energy under fluctuating light conditions. Many of the core

reactions and features of the PSs are highly conserved among photosynthetic organisms. Research on the structure and function of the core components of the photosystems and thylakoid membranes often uses phototrophic microorganisms for the ease of maintaining cultures and isolating or purifying components. These include the green alga *Chlamydomonas reinhardtii*, diatoms such as *Phaeodactylum* and *Thalassiosira*, and cyanobacteria such as *Synechocystis* and *Synechococcus*. With microorganisms, it is also possible to study in vivo and whole organism properties of photosynthesis without some of the many confounding processes present in land plants, such as stomatal conductance of CO<sub>2</sub>, water relations, and carbohydrate allocation (source–sink) dynamics.

A lot of photosynthesis research today is directed toward improving photosynthesis to create more climate-resilient crops and enhance crop yields in the face of global climate change (see the companion article in this issue by [Croce et al. 2024](#)). And yet, despite the long history of photosynthesis research (see e.g. [Nelson and Junge 2015](#); [Stirbet et al. 2020](#); [Sharkey 2022](#)) and significant advances we have seen in recent years (e.g. [Croce and van Amerongen 2020](#); [He et al. 2020](#); [Perera-Castro and Flexas 2020](#);

Received January 23, 2024. Accepted July 15, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of American Society of Plant Biologists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Bhowmick et al. 2023; Greife et al. 2023), we do not understand many aspects of photosynthetic energy conversion, such as details of its evolution, dynamics, and regulation.

In this commentary, a number of researchers working on fundamental aspects of photosynthesis pose and discuss what they view as the most compelling open questions in their areas of research. We begin at the reaction center and catalytic environment of the Mn cluster and move outwards to address questions related to light harvesting, photosystem evolution, thylakoid membrane dynamics, and alternative electron flow pathways. We have avoided questions related to Rubisco, as these have been covered in many recent reviews and perspectives (Carmo-Silva and Sharwood 2023, and references cited therein; Croce et al. 2024). Two final sections address the link between photorespiration and stomatal control and adaptive changes that make an enzyme suitable for its function in the  $C_4$  carbon-concentrating pathway present in important crop species.


## How does the catalytic environment of the $Mn_4CaO_5$ cluster accommodate and control the multielectron/multiproton reaction during water oxidation?

(Written by Vittal K. Yachandra, Jan Kern, and Junko Yano)

The model for photosynthetic water splitting proposed by Kok et al. (1970), based on the yield of dioxygen as a function of light flashes, has served as a blueprint for all subsequent work on the mechanism of the water-splitting reaction. This kinetics-based model proposed the advancement of the oxygen-evolving

complex (OEC), the site where the oxidation of water occurs in photosystem II (PSII), through five intermediates known as the S-states ( $S_i$ ,  $i=0$  to 4). The absorption of four photons, one at a time, advances the OEC from  $S_0$  through  $S_4$ , after which  $O_2$  is released. This model, called the Kok S-state cycle or clock (Fig. 1A), provided an elegant explanation for the coupling of the one-electron photochemistry occurring at the PSII reaction center to the four-electron redox chemistry occurring at the OEC. The working hypothesis was that upon the absorption of a photon, an oxidation equivalent is stored on the OEC, and after reaching the  $S_4$  state, the most oxidized intermediate, two water molecules are oxidized to form one molecule of  $O_2$ . The  $S_0$ ,  $S_1$ ,  $S_2$ , and  $S_3$  states are stable and can be trapped, but the  $S_4$  state is normally referred to as a short-lived or transient state that appears during the  $S_3$  to  $S_0$  transition prior to O–O bond formation and release of  $O_2$ . The Kok cycle also entails the release of four protons and four electrons. The consensus is that the protons are released in a 1, 0, 1, 2 pattern for the  $S_0$  to  $S_1$ ,  $S_1$  to  $S_2$ ,  $S_2$  to  $S_3$ , and  $S_3$  to [ $S_4$ ] to  $S_0$  transitions, and the electrons extracted from the donor OEC site by the PSII reaction center are transported via the electron transfer components to the quinones on the acceptor side.

Identifying the structure of PSII and the Mn-containing catalytic metal cluster of these intermediates, especially, during the  $S_3$ –[ $S_4$ ]– $S_0$  transition has been challenging and the subject of many studies (reviewed in (Wydrzynski and Satoh 2005; Nelson and Yocum 2006)) using diverse spectroscopic techniques, such as X-ray (Yachandra et al. 1996; Dau and Haumann 2008; Yano and Yachandra 2014), Fourier Transform Infrared Spectroscopy (FTIR; Debus 2015; Nagao et al. 2018), electron paramagnetic resonance (EPR; Lubitz et al. 2019), and X-ray crystallography



**Figure 1.** The Kok cycle and structure of PSII. **A)** The Kok cycle is shown in the middle, with the structures from XFEL crystallography of the S-states and at intermediate time-points between the  $S_2$  and  $S_3$ ,  $S_3$  and  $S_0$  states. O5 is shown in blue, and in red is the insertion of OX between Mn1 and Ca during  $S_3$  to  $S_0$  and its disappearance between the  $S_3$  and  $S_0$  transitions. The sigma value in crystallography is used for the scaling of the electron density; a higher value shows higher accuracy. **B)** On top is the structure of PSII, showing the electron transfer components, the OEC, and the proton and water channels (in red, blue, and green) in one monomer (left). The transmembrane helices and the helices in the extrinsic polypeptides are shown in color in the monomer on the right. Below this is the oxo-bridged Mn<sub>4</sub>Ca cluster in the  $S_1$  state in detail, with the water and other terminal ligands of Mn and Ca provided by the protein glutamate, aspartate, and histidine residues.

(Umena et al. 2011). These studies revealed that the catalytic metal site consists of an oxo-bridged Mn<sub>4</sub>Ca hetero-nuclear cluster (Fig. 1B) that accumulates charge as it proceeds through the S-state cycle, reinforcing the hypothesis that the S<sub>0</sub> through S<sub>3</sub> transitions are primarily where the oxidizing equivalents are stored, with little or no specific chemistry related to the O–O formation from two water molecules. However, it was shown that during progression from S<sub>0</sub> through the S<sub>3</sub> state, the oxidation equivalents or the charge is not centered only on the Mn but more delocalized onto the ligands (Glatzel et al. 2004, 2013). There was no evidence for the presence of a peroxo or other intermediates, which are intermediate species one can expect for a water oxidation reaction. Intermediates such as metal-peroxo, -oxo, and -superoxo species have been detected in the study of inorganic water-oxidation catalytic systems. These data indicated that the entire chemistry involving water and the O–O bond formation occurs during the last step. This is the discharge/reduction step, where the most oxidized S<sub>4</sub> state is reduced to S<sub>0</sub> state (most reduced) with concomitant oxidation of water either in one fell swoop or in a stepwise manner, in which case it would proceed through the expected intermediates of water oxidation, with the most likely being two two-electron steps resulting in dioxygen via a transient peroxo intermediate.

The recent advent of X-ray free-electron laser (XFEL)-based room temperature crystallography and X-ray spectroscopy has had a profound influence on the structural studies and on the study of the mechanism of the water-splitting reaction (Kern et al. 2012, 2013, 2014; Suga et al. 2015). The structures of the S<sub>0</sub>, S<sub>1</sub>, S<sub>2</sub>, and S<sub>3</sub> states have been determined at ~2 Å resolution (Fig. 1B). Furthermore, it is now possible to follow the structural changes between the S<sub>2</sub> and S<sub>3</sub> states (Ibrahim et al. 2020) where the insertion of a new water occurs between Mn1 and Ca (likely as –OH after releasing one proton), and most importantly between the S<sub>3</sub> and S<sub>0</sub> states where the accumulated oxidizing equivalents are reduced and O<sub>2</sub> is released (Fig. 1B) (Bhowmick et al. 2023).

Several events accompany the S<sub>3</sub> to S<sub>0</sub> transition that resets the Kok-clock to the S<sub>0</sub> state, which includes the release of two protons, the release of dioxygen, and the recovery of one water at the OEC. The newly introduced ligand between Mn and Ca in the S<sub>3</sub> state disappears during the S<sub>3</sub> to S<sub>0</sub> step, and this is related to the release of O<sub>2</sub>. The release of two protons signaled by changes in the gate region in the proton channel (Hussein et al. 2021) have been observed at two different time-points in the XFEL crystallography data (Bhowmick et al. 2023). Yet, it is not clear how exactly O–O formation and the recovery of the catalytic center takes place, due to the resolution of the current crystallography data that hinders accurate modeling of oxygen positions in the fraction of the changing population. The reduction process may create a longer lived intermediate, perhaps a peroxo species, which is two-electron reduced from the most oxidized S<sub>4</sub> state. This is one step away from a further two-electron reduction of the OEC that results in the release of O<sub>2</sub>. We await structural studies at higher resolution.

The ability to visualize PSII structural changes through the catalytic process and understand the importance of the overall protein structure in the water-oxidation mechanism has also opened up new questions that are perhaps foundational to many multielectron catalytic reactions. How does the protein environment modulate the structure to neutralize the charge density changes on the OEC during catalysis? How does the water network, together with the amino-acid residues, facilitate proton transfer and substrate transport? How are the reaction kinetics affected by different environmental parameters such as pH and

temperature? And how does the fundamental mechanism of the OEC remain the same through the diverse evolutionary tree (Hussein et al. 2023)? Recent advances in cryogenic electron microscopy in combination with site-specific mutations will likely contribute to answering some of these questions. While *Thermosynechococcus vestitus* (previously known as *T. elongatus*) has often been used for crystallography due to the availability of high-quality crystals, more species are being used for structural studies with cryogenic electron microscopy, and the structures from different organisms have been compared. Such comparisons can help to address questions about the fundamentally important design components for the water oxidation reaction in nature. In addition, the above-mentioned XFEL crystallography at room temperature, along with various spectroscopic methods, enables us to bridge structural information with spectroscopic data and provide the spatial and temporal sequence of events during the reaction. Thus, the study of the light-driven water oxidation reaction in PSII provides a unique opportunity to understand how the metal catalytic center and protein environment orchestrate and enable the multielectron/multiproton reaction, providing fundamental knowledge that goes beyond the water oxidation reaction.

## How do protein networks control light harvesting?


(Written by Gabriela Schlau-Cohen and Dvir Harris)

In photosynthetic light harvesting, a network of chromophore-containing antenna proteins absorbs light and transports photoenergy over distances of tens to hundreds of nanometers to reach reaction centers (Fig. 2). Despite the long distances, this transport can occur with near-unity quantum efficiency. At the same time, the efficiency is actively regulated by the network in a photoprotective response to light levels (Eberhard et al. 2008). The functionality of the network relies on the collective behavior of the constituent proteins. Yet studies have typically been limited to individual proteins, where relevant properties are often absent, or intact systems, where contributions from many proteins are combined to obscure behaviors of interest (van Grondelle and Novoderezhkin 2006). Resolution of the interactions and dynamics between proteins has been a long-standing challenge, leading to the major open question: *How do protein networks control light harvesting?* We discuss three facets of this open question.

## How is energy transferred between proteins?

The efficient transport of photoenergy through the antenna complexes to the reaction center is well established in photosynthetic organisms. Such long-distance transport emerges from a series of energy transfer steps within photosynthetic membranes. Cumulatively, these steps contribute to the hundreds of picoseconds timescale of solar energy conversion, whereas the relaxation time within individual proteins is approximately only one picosecond. The vast majority of previous experiments focused on individual proteins, even though transfer between proteins occupies most of the total time and is the key to long-distance energy transport (Fiebig et al. 2023). Despite its crucial importance, protein-to-protein energy transfer has been challenging to measure.

Several questions remain about how protein-to-protein energy transfer enables robust long-distance transport. In green plants, the energetic gradient of the protein network towards the reaction



**Figure 2.** PSII from green plants. Chromophore-containing antennae capture and transport light energy (orange) to the PSII reaction center, where charge separation occurs (black). Under high light conditions, the local cellular environment and the organization of the protein network activate photoprotection within the antenna proteins, likely involving a switch of the photophysics of the embedded chlorophyll and carotenoids associated with different structural states (PDB: 5XNM). The protein network is formed through van der Waals interactions between proteins.

enter is nearly flat (Mirkovic et al. 2017). How a nearly flat gradient produces an effective directional energy flow remains unclear. Both the composition and organization of the protein network also change under light conditions. Energy transfer is highly sensitive to distance, and so even distance changes of a single nanometer can change the rate by an order of magnitude. Despite this sensitivity, the quantum efficiency of light harvesting changes minimally—less than 10%—with network organization (Fleming et al. 2012). Thus, how network reorganization impacts energy transport is not fully understood.

Despite the inherent challenges, several promising directions can now probe protein-to-protein energy transfer. Because of spectroscopic improvements, energy transfer rates have been extracted for spectrally shifted antenna proteins in membrane fragments or in vivo (Tiwari et al. 2018; Lüer et al. 2012). Alternatively, model membrane systems have been used to biochemically isolate antenna proteins for measurements of protein-to-protein energy transfer (Dai et al. 2018; Wang et al. 2023a, 2023b, 2023c). Finally, an integrative approach using single-molecule spectroscopy and cryogenic electron microscopy probed the heterogeneity associated with antenna to reaction center energy transfer (Harris et al. 2023). These recent technological advances enable investigation of the important and fundamental process of energy transfer between proteins.

### How does the network regulate photoprotection?

In oxygenic photosynthesis, excess energy can cause photooxidative damage to proteins. Under high light conditions, a drop in the luminal pH activates a series of photoprotective processes, known as nonphotochemical quenching (NPQ). The fastest NPQ process is qE, the dissipation of excess energy as heat. The major antenna protein, light-harvesting complex II (LHCII), is a major site of dissipation for PSII (Schlau-Cohen et al. 2015; Nicol and Croce 2021), and its organization within the protein network changes, notably through the formation of clusters (Goral et al. 2012; Ruban and Johnson 2015). Recapitulating the physiological

LHCII–LHCII interactions responsible for the functional switch has remained challenging, and so how these interactions regulate the dissipative pathways of photoprotection in vivo remains unclear.

Studies on aggregates and arrays of LHCII, which aim to emulate LHCII clustering, have shown enhanced dissipation, and a pH dependence reminiscent of the in vivo conditions (Barros and Kühlbrandt 2009; Petrou et al. 2014; Son et al. 2021). A dissipative photophysical pathway, chlorophyll-to-carotenoid energy transfer, was recently clearly resolved with the cluster and pH dependence expected for photoprotection. In recent in vivo measurements, the carotenoid photophysics correlated with NPQ (Park et al. 2019), suggesting that the photoprotective switch involves these important biomolecules. Such effects may be induced by LHCII clustering.

### What are the protein–protein interactions that govern network architecture?

In recent years, the organization of the protein networks has been revealed in exquisite detail (Wietrzynski et al. 2020; Opatíková et al. 2023; Feng et al. 2023). Advances in structural biology, including electron microscopy and tomography at cryogenic temperatures and atomic force microscopy, have provided snapshots that dramatically advanced our understanding. However, the interactions between the proteins that drive the formation of these networks and their reorganization in response to environmental stimuli remain unknown. While a small number of computational studies have investigated these interaction energies (Schneider and Geissler 2013; Mao et al. 2023), the well-established deviation between computational energies and experimental benchmarks (Keskin et al. 2016) means that new methods are needed to determine the thermodynamic driving forces behind protein networks. Recent experimental measurements of the interaction energies between LHCII revealed a net attraction of  $\sim -5 k_B T$ , which represents an enthalpically dominated thermodynamic driving force behind LHCII clustering. Such studies capture the interactions

behind the membrane organization of PSII from the perspective of equilibrium statistical thermodynamics, which has a long and rich tradition in biology (Manna et al. 2023).

In conclusion, structural and spectroscopic advances over the past decades built an understanding of the function and composition of the protein networks of photosynthetic light harvesting. In particular, the structure, organization, and dynamics of the constituent proteins have been revealed. The open questions sit at the interface between these proteins. Resolution of the interactions between proteins and how these interactions modulate function requires further improvements to our tools that characterize and manipulate these biosystems. Answering the central question—How do protein networks control light harvesting?—remains a grand challenge for the field.

## Why do diatoms have so many different light-harvesting proteins?

(Written by Nicolas Herrmann and Claudia Büchel)


Vascular plants express an array of light-harvesting complexes (LHC) and the different functions of these LHCS as major or minor antennae for PSII, as PSI-specific antennae, or as mobile antennae in the context of state transitions are well described (Croce and van Amerongen 2020). Algae, by contrast, contain many more Lhc proteins (Lhc and LHC typically are used to denote the algal and vascular plant proteins, respectively), and one of the intriguing questions is the reason behind this diversity. Besides the well-studied green algae, there are many algal groups that use different pigments and even different proteins for light harvesting (Büchel 2015). Most prominent are the Stramenopiles, which are distantly related to green algae and vascular plants. These algae are derived from a so-called secondary endosymbiosis, where a eukaryotic host engulfed an organism related to red algae that was evolutionarily reduced to the chloroplast (Bhattacharya and Medlin 1995). The group of Stramenopiles with the best-studied light-harvesting systems are diatoms, unicellular organisms that are responsible for 20% of primary production worldwide (Field et al. 1998). Diatoms are found in many different habitats, ranging from biofilms to free-living marine or limnic planktonic forms. Marine diatoms are relatively easy to culture, and after whole-genome sequences as well as molecular tools for transformation became available, different species emerged as reference systems, i.e. the pennate diatom

*Phaeodactylum tricornutum* and the centric diatoms *Thalassiosira pseudonana*, *Cyclotella meneghiniana*, and *Chaetoceros gracilis*.

Diatoms possess a photosynthetic apparatus that is similar to that of vascular plants, including membrane-intrinsic antenna proteins of the Lhc family (Büchel 2015). However, their pigmentation is very different. Whereas chlorophyll (Chl) *a* is present in diatoms, Chl *b* is missing and only 2 to 3 molecules of Chl *c* are bound instead. The carotenoid fucoxanthin (Fx) is the main accessory pigment with up to seven molecules per protein of the eponymous Fucoxanthin-Chlorophyll Proteins (FCP). The Fx molecules are bound to various sites of FCP, inducing differences in absorption (Fig. 3A). Thus, FCP absorbs up to 585 nm, giving rise to the characteristic brown color of diatoms and optimizing photosynthesis in deeper water, where blue-green light dominates.

A wealth of spectroscopic data exists about the excitation energy transfer inside FCP complexes. Fx displays efficient excitation energy transfer to Chl *a* (Papagiannakis et al. 2005). The fastest transfer (~30 to 60 fs) takes place between Chl *c* and Chl *a* (Butkus et al. 2015). Thus, excitation energy transfer between different Chl molecules is faster and FCP rely more on carotenoid absorption for light-harvesting than do Lhc of vascular plants. Nonetheless, these features are similar for all Lhc proteins from diatoms studied so far and do not shed light on a possible functional heterogeneity.

All diatom species contain a much higher number of expressed Lhc genes than found in vascular plants. *T. pseudonana* has more than 30 FCP proteins, *C. meneghiniana* has 23 FCPs, and *C. gracilis* and *P. tricornutum* have 22 and 32 FCPs, respectively (Kumazawa et al. 2022). PSI of diatoms is surrounded by monomeric FCPs, but the antenna is much larger than that of vascular plants, comprising ~16 to 24 different FCPs in the case of *C. gracilis* (Nagao et al. 2020; Xu et al. 2020) (Fig. 3B). PSII structures are available from three centric species that differ tremendously in their FCP arrangement and composition, a diversity completely unknown in vascular plants. Whereas *C. gracilis* has three monomeric and two tetrameric FCP per core monomer (Pi et al. 2019), PSII of *C. meneghiniana* (Wang et al. 2023a, 2023b, 2023c) has only six FCP proteins. The high-resolution structure of *T. pseudonana* PSII shows one additional FCP (Feng et al. 2023) and is in contrast to the low-resolution structures, where three monomers and one trimer were modeled (Arshad et al. 2021). In addition, even the oligomeric state of FCP complexes found in the free pool of FCPs differs between species: for *P. tricornutum* Lhc4 homodimers (Wang et al. 2019) and different trimers (consisting of the other FCP



**Figure 3.** The FCP complex. **A)** Absorption spectrum of an FCP complex (black, FCPs from *C. meneghiniana*) and its pigments. Fx molecules are bound at different sites, absorbing more to the red wavelength range (Fx “red”), intermediate (Fx “green”), or to the blue (Fx “blue”). **B)** Scheme of FCP arrangement in PSI, PSII, and the free FCP pool of different species. Lhc proteins are shown in red, other FCPs are colored according to their subfamily (light purple, Lhcf; blue, Lhcr; purple, lhcg; gray, unknown). For references, see text.

proteins including Lhcf4) (Gundermann et al. 2013) were reported, whereas analysis of *C. meneghiniana* revealed trimers and nonamers (Röding et al. 2018; Wang et al. 2023a, 2023b, 2023c). Oligomerization influences the pigment network and correspondingly, excitation energy transfer into the reaction centers. So far, no functional reason is known for this huge inter-species diversity.

To cope with rapid changes in light intensity while maintaining optimum photosynthetic energy flux, diatoms employ nonphotochemical quenching (NPQ), which is very fast and has a high capacity compared to NPQ of vascular plants (Ruban et al. 2004). NPQ is triggered by a high trans-thylakoid pH gradient under strong light and relies on a xanthophyll cycle that converts diadinoxanthin (Ddx) to diatoxanthin (Dtx) under excess light (Lavaud et al. 2002) and special FCP proteins called Lhcx. Diatoms contain many Lhcx proteins. Lhcx1 was demonstrated to be involved in NPQ in *P. tricornutum* (Bailleul et al. 2010) as well as in *C. meneghiniana* (Ghazaryan et al. 2016), but Lhcx1 cannot sense pH (Buck et al. 2021) and therefore the mechanism triggering NPQ remains enigmatic. Aggregation of FCPs containing Lhcx1 from the *C. meneghiniana* FCP pool reduced fluorescence in vitro and Dtx enhanced this fluorescence, which resembles NPQ in vivo (Gundermann and Büchel 2012). Domain swap experiments demonstrated the importance of a Trp residue of Lhcx1 that is close to the presumed Ddx binding site (Buck et al. 2021), hinting at a direct connection between Lhcx proteins and Ddx/Dtx. However, although the participation of Dtx in NPQ is well established, an in-depth spectroscopic analysis will be required to prove that Dtx that is bound to Lhcx1 participates in NPQ. The centric diatoms *T. pseudonana* and *C. meneghiniana* contain a special Lhcx protein, Lhcx6\_1, which was found to be located in PSII complexes, harbors Dtx, and might be the second quenching site usually found in centric diatoms (Calvaruso et al. 2020, Gruneva et al. 2008). The sole presence of Dtx in an Lhcx structure does not prove its involvement in NPQ. Various abiotic stresses were shown to regulate the expression of the other three Lhcx proteins in *P. tricornutum* (Taddei et al. 2016), tuning NPQ. This functional diversification might help diatoms cope with variable environments.

Our understanding of the arrangement of FCPs around the photosystems, pigment organization, and excitation energy transfer has greatly improved in recent years. But why is FCP organization around PSII so diverse among the different species and what are the roles of the different FCP proteins? Involvement of FCPs in regulatory functions is the obvious idea to explain the huge diversity in and between species, but whether this explanation is true remains mostly enigmatic, as plasticity in the spatial arrangement of different FCPs remains unreported. And how precisely are the different Lhcx proteins involved in NPQ? Answering these questions will help us to understand the huge ecological and evolutionary success of diatoms.

## New perspectives on the evolution of the photosystems: Are the photosystems evolvable?


(Written by Tom Dongmin Kim, Emma Chaloner, and Tanai Cardona)

How and when photosynthesis originated remain compelling questions in the Life and Earth Sciences. How photosystems, the multifactor and multiprotein assemblies that power primary production, were put together from their constitutive parts and for what original function, are questions that still need answers.

The photosystems are well over three billion years old and are potentially amongst the oldest enzymes (Oliver et al. 2023). On the one hand, that such endurance is possible suggests that photosystems have an in-built capacity for evolution and adaptability that enable photochemical processes to occur in as varied and dynamic conditions as can be found on Earth's photic zones, and on a planet that has gone through remarkable transformations over the eons. On the other hand, photosystems are highly conserved and slowly evolving enzymes, often perceived as somewhat immutable, even dubbed "frozen metabolic accidents" (Shi et al. 2005). Given that all photosystems share a common origin, evidence for the photosystems' evolvability is most conspicuously noted when comparing PSII and its unparalleled capacity to split water by generating some of the most oxidizing species in biology, with PSI, which has evolved to generate some of the most reducing ones.

These evolutionary considerations trigger a question that has received little explicit attention in the field of photosynthesis research: *are the photosystems evolvable?* Given that photosystems are biological systems, it is safe to assume that they are. However, the question raises a number of additional fundamental considerations about how photosystems have changed through time. For example, what are the molecular mechanisms that make the photosystems adaptable? How quickly can a photosystem gain a new function given adequate selective pressure? And, given selective pressures not found in nature, what new photochemical and catalytic properties could a photosystem evolve? We could go one step further: *Can photosystems be evolved in the lab?* We propose here that in an endeavor to answer these questions, we could gain new insights into why the photosystems evolved to be the way they are and open new pathways to greener chemical and biotechnological processes.

Photosystems are modular and this modularity is the basis for their capacity to remain adaptable. In the wild, this means that the photochemical or catalytic properties of a photosystem can be modified to respond to the environment by exchanging subunits, which work as replaceable modules. The better-known example of this is the fine-tuning of PSII energetics to perform optimally under different light intensities, and the most extreme example is the transformation of PSII from a water-oxidizing to a Chl *f*-producing enzyme (reviewed recently by Oliver et al. (2023)). In both cases, these functional alterations are achieved by replacing a standard D1 subunit for a variant form of D1. And in both cases, PSII evolved new photochemical or catalytic properties by acquiring modifications on a spare copy of a gene encoding the D1 subunit within the genome of cyanobacteria. Alterations of PSI by a similar mechanism are also known. For example, some cyanophages carry a set of PSI genes that are deployed upon infection of the cyanobacterium. In this adaptation, the standard PsaF and PsaJ subunits are replaced by a subunit encoded in the cyanophage's genome which is a fusion of the two (Sharon et al. 2009). This novel subunit has features that make the complex more promiscuous to protein electron carriers (Mazor et al. 2014). Another adaptation commonly observed in heterocystous cyanobacteria and their close relatives is the exchange of PsaB paralogs, which is likely to lead to a fine-tuning of the energetics of the complex in ways and conditions that are yet to be understood (Gisriell et al. 2023). Some strains encode up to four distinct PsaB versions. While the exchange of just a single subunit is sufficient to modify the photochemical and catalytic properties of the photosystems, there is no appreciable boundary as to the extent of change that can occur driven by sustained evolutionary pressures. A clear example of this is the far-red light photoacclimation response



**Figure 4.** Conceptualization of evolved photosystems. **A)** A PSII evolved to bind a dinuclear Ru cluster ( $\text{Ru}_2\text{O}_2$ ), which catalyzes a hypothetical specialized oxidative reaction. **B)** A hypothetical chimeric PSI fused with a reductive dehalogenase and evolved for optimal electron transfer to the catalytic site at a cobalamin cofactor ( $\text{B}_{12}$ ). Protein scaffolds of the photosystems are shown in transparent gray ribbons, and that of the reductive dehalogenase in magenta; chlorophylls are shown in green, carotenoids in purple, and iron–sulfur clusters are depicted with yellow and orange spheres. PSII structure was modified from PDB ID 3WU2, PSI from 1JB0, and reductive dehalogenase from 5M2G.

(Gan et al. 2014), which involves the swap of at least five subunits of PSII, and six of PSI, and the use of three distinct chlorophyll types to enable oxygenic photosynthesis under far-red light and in the absence of visible light.

The next step towards understanding the evolution of the photosystems is to evolve them in a laboratory setting and attempt to change their catalytic and photochemical properties purposefully. There is experimental evidence suggesting that the modularity of PSII can facilitate the application of directed evolution methods. For example, random mutagenesis of D1 was used to select for PSII complexes more tolerant to high light intensities (Narusaka et al. 1999), with tolerance to herbicides (Narusaka et al. 1998), and with tolerance to ionizing radiation (Rea et al. 2011). However, these studies stopped at a single round of mutagenesis and selection. In addition, directed evolution was used to successfully change the directionality of electron transfer of the purple bacteria reaction center (Faries et al. 2012).

Evolving PSII could involve modifications of the electron donor side to change the nature of the water-oxidizing cluster and to enable access to other substrates (Fig. 4A). For example, to study one of the plausible transitional stages in the origin of PSII (Chernev et al. 2020), it may be possible to evolve a PSII that can efficiently oxidize Mn but cannot split water. Alternatively, a PSII could be evolved to decompose a complex organic pollutant in a sequential multielectron multiproton extraction process. The energetics of charge separation could be optimized for the specific catalytic demands imposed by the new substrates. There is evidence supporting the idea that  $\text{P}_{\text{D1}}$ , one of the key redox chlorophylls of PSII that has a midpoint potential of about +1.2 V, is naturally tuned down by  $-0.14$  V to compensate for electrostatic effects created by the  $\text{Mn}_4\text{CaO}_5$  cluster (Ishikita et al. 2006), implying that the photochemical core could be pushed to generate even higher oxidizing potentials if desired. Evolving PSI could involve adding novel catalytic domains, which could be further optimized with directed evolution methods. In this sense, PSI could be engineered to transfer electrons from the terminal iron–sulfur clusters to the cofactors of the new domains, resulting in novel photobio catalysis (Fig. 4B). This concept has been demonstrated in the genetic fusion of the extrinsic subunits of PSI with hydrogenase enzymes (Appel et al. 2020; Kanygin et al. 2020; Wang et al. 2023a, 2023b,

2023c). The chimeras were all characterized as functional photosystems capable of light-driven  $\text{H}_2$  evolution. Such experiments, while opening new avenues for applications, would also test many existing rationales aimed at explaining photosystem structure and function. For example, what are the exact energetic tradeoffs between photochemical efficiency and photoprotection? Thus, if oxygen were not to be a reaction by-product of a hypothetical evolved PSII oxidizing a new substrate, could energy that is sacrificed by the reaction center to prevent back-reactions be redirected towards improving turnover efficiency? If that were the case, then, what would be the maximum rate of water oxidation of PSII, i.e. if the system had not needed to evolve into its bioenergetics protective mechanisms against the production of reactive oxygen species? By exploring these questions, a positive feedback loop can be created between research aimed at understanding photosystem evolution and developing new photosystem-based technologies for better biocatalysis.

To conclude, two specific questions can be asked to guide new research. First, what are the limits of photosystem evolvability? The answer to this question will help us understand the limits of light-dependent life on Earth and extrasolar planets in habitable zones. It can also help us understand why life came to be the way it is. Second, how can novel light-driven enzymes, featuring customized photochemical and catalytic properties, contribute to making the chemical and biotechnological industries more sustainable? This question is an invitation to a new generation of photosynthesis researchers to use their creativity in exploring how new photosystems could be harnessed as solutions to some of the global challenges we face.

## Thylakoid membrane dynamics: How are protein biogenesis and mobility orchestrated in this densely packed system?

(Written by Conrad Mullineaux)

Thylakoid membranes in chloroplasts and cyanobacteria are densely packed with protein complexes. They present a crowded environment, not only in the plane of the membrane but also in the third dimension, where parallel membrane surfaces may be

tightly appressed, or in close proximity and sandwiching a dense mass of soluble protein. Dense packing of the system allows a high concentration of pigment-protein complexes, which is necessary for the efficient absorption of sunlight. However, it creates challenges for other aspects of membrane function, including regulatory re-organization, diffusion of mobile electron carriers, and the biogenesis of protein complexes, which could all be impeded by the crowded environment in and around the thylakoids (Kirchhoff 2014a).


The translation and membrane insertion of membrane-integral protein complexes invariably seem to occur in specialized regions of the membrane that may be quite remote from much of the mature functioning thylakoid surface. This likely arises because the sheer bulk of the ribosomes excludes them from most of the crowded thylakoid system. In cyanobacteria, ribosomes are found only at the proximal thylakoid surface adjacent to the central cytoplasm and are therefore excluded from about 90% of the thylakoid system (Fig. 5; Rast et al. 2019; Mahbub and Mullineaux 2023). mRNAs encoding membrane-integral thylakoid proteins are found at the same membrane surface (Mahbub et al. 2020; Mahbub and Mullineaux 2023). In chloroplasts of the green alga *Chlamydomonas*, translation is heavily focused on a specialized “T-zone” surrounding the pyrenoid (Schottkowski et al. 2012). In vascular plant chloroplasts, ribosomes are excluded from the tightly appressed grana membrane surfaces, but ribosomes also appear very scarce across large areas of the stromal lamellae, being concentrated at the most peripheral regions of the thylakoid that have unimpeded access to the chloroplast stroma (Shimoni et al. 2005).

If translation at the thylakoid is highly localized, it raises the question of how newly synthesized protein complexes can find their place in the mature thylakoid membrane. Thylakoid membranes seem to be more interconnected than is often apparent at first sight. For example, the thylakoids of the cyanobacterium *Synechococcus elongatus* approximate a set of nested cylinders (Fig. 5), but electron tomography revealed membrane bridges that interconnect the whole system (Nevo et al. 2007). So, it is likely that there is a continuous membrane surface connecting every region of a thylakoid to a translation zone, but the problem

of restricted protein diffusion in the crowded thylakoid membrane environment remains—is it feasible for a newly synthesized complex to diffuse from the site of its translation to the furthest parts of the thylakoid system? Taking *S. elongatus* as an example, Fluorescence Recovery after Photobleaching (FRAP) measurements on GFP-tagged photosynthetic complexes suggest long-range lateral diffusion coefficients of the order of  $10^{-10}$   $\text{cm}^2\text{s}^{-1}$  (Casella et al. 2017). Such a diffusion coefficient would permit a complex to diffuse on average 1  $\mu\text{m}$  from its starting point in 25 s, however for radial diffusion from the proximal to the distal layers of the thylakoid system the problem is hugely compounded by the need for the complex to encounter one of the very occasional bridges that connect the concentric thylakoid membrane layers (Nevo et al. 2007). There is an analogous problem in chloroplast appressed grana membranes, where FRAP measurements suggest a limited mobile pool of chlorophyll-protein complexes with a diffusion coefficient of about  $5 \times 10^{-11}$   $\text{cm}^2\text{s}^{-1}$  (Kirchhoff et al. 2008), but there are restricted connections between the appressed membranes and the stromal lamellae (Mustardy et al. 2008). Computational modeling combined with high-resolution data on 3D thylakoid ultrastructure will be needed to define the extent of the problem.

If, as seems likely, movement of newly synthesized complexes to the furthest parts of the thylakoid system is slow, then what are the solutions and what are the functional implications? Taking *S. elongatus* (Fig. 5) as an example, two extreme scenarios could be envisaged for the incorporation of new complexes into the expanding thylakoid membrane as the cell grows. In the first scenario, new complexes stay close to their site of translation. As the cell grows and more complexes are synthesized, the parts of the proximal membrane surface mature into fully functional thylakoid. New areas of biogenic membrane surface are generated to replace them, so that new thylakoid is created at the proximal side of the thylakoid system and older areas of thylakoid are gradually pushed towards the cell periphery as the cell grows. One piece of evidence against this idea is that in *S. elongatus* cells that are regenerating their thylakoid system after growth in high light, new thylakoid membrane sacs seem to appear at the distal edge of the system between the existing thylakoids and the plasma membrane (Huokko et al. 2021). These new membranes must presumably be populated by protein complexes migrating from the proximal side of the system. The second scenario assumes that the proximal thylakoid membrane surface remains as a biogenic surface throughout, and complexes migrate out of it as they are assembled. It seems that this idea must apply at least to the replacement of the D1 subunit of PSII, which is rapidly turned over in high light (Komenda et al. 2012). Newly translated D1 proteins must surely be needed for repair of damaged PSII complexes throughout the thylakoid system. Interestingly, there are indications that the mobility of chlorophyll-protein complexes increases following high light exposure, in both chloroplasts (Goral et al. 2010) and cyanobacteria (Sarcina et al. 2006).

Both scenarios outlined above would predict gradients in thylakoid membrane protein composition depending on their distance from the nearest translation site. Repair and regulatory adjustments to the population of photosystems (e.g. (Dietzel et al. 2008; MacGregor-Chatwin et al. 2022)) might take effect at different speeds in different regions of the membrane, with strong functional consequences. Understanding the consequences will require progress in cell biological approaches to thylakoid biogenesis, to complement the remarkable advances from biochemical and structural approaches (e.g. (Zabret et al. 2021)). Electron tomography has provided new insight into the 3D architecture of



**Figure 5.** The photosynthetic protein biogenesis problem, illustrated by a schematic cross-section of a cell of the cyanobacterium *Synechococcus elongatus*. Translation of membrane-integral photosynthetic proteins occurs only at the proximal thylakoid surface facing the central cytoplasm and the nucleoid. How do new complexes get to the more distant parts of the thylakoid system?

thylakoids, as in (Shimoni et al. 2005; Nevo et al. 2007; Engel et al. 2015; Rast et al. 2019), and this should be coupled with fluorescence microscopy to probe membrane dynamics. Techniques such as structured illumination microscopy, as in (Wood et al. 2018), will allow higher resolution than was previously achievable. It would be exciting to develop fluorescent labeling methods that distinguish between newer and older photosystems. This could reveal the flux of the membrane system over time, as the cell or organelle expands and divides. How do the dynamics of the membrane and the locations of membrane protein production constrain photosynthetic function, and how much do they restrict acclimation to new conditions? These will be overarching research questions for the next years.

## How fluid are thylakoid membranes?

(Written by Wojciech Wietrzynski and Benjamin D. Engel)

One of the most intriguing characteristics of thylakoid membranes is their diverse molecular architecture (Perez-Boerema et al. 2024). In cyanobacteria, as well as the chloroplasts of red algae and glaucophytes, thylakoids are decorated with bulky phycobilisome light-harvesting antennae, which space the membranes apart and prevent stacking interactions. Here, PSII has been observed to form linear arrays, bound underneath rows of phycobilisomes (Rast et al. 2019; Li et al. 2021a, 2021b). PSI has been observed to bind alongside these PSII arrays in red algae (You et al. 2023), or in cyanobacteria to form trimers or tetramers occupying separate domains within the nonstacked thylakoids (MacGregor-Chatwin et al. 2017).

In contrast, thylakoid architecture in the chloroplasts of green algae and plants can generally be subdivided into appressed and nonappressed (stacked and unstacked) domains (Mustárdy et al. 2008). These chloroplasts lack phycobilisomes and instead rely on light-harvesting complexes (LHCs) and related membrane-integral antenna proteins, which do not space the thylakoid membranes apart. In fact, there is evidence that interactions between LHCII proteins contribute to membrane stacking in plants (Standfuss et al. 2005; Daum et al. 2010; Zeno et al. 2022). Stacking enforces lateral heterogeneity in thylakoid membranes: PSI and ATP synthase populate the nonappressed domains, whereas PSII is restricted to the appressed domains, and only cytochrome *b<sub>6</sub>f* is abundant in both domains (Andersson and Anderson 1980; Wietrzynski et al. 2020). This segregation of components prevents energy spillover from PSII to PSI, and has deep implications for the regulation of photosynthesis. Balancing of PSI and PSII excitation via migration of LHCII between domains (state transitions; Allen and Forsberg 2001), repair of PSII complexes damaged by photoinhibition (Barber and Andersson 1992), and integration of newly synthesized proteins into the appressed membranes would all require the exchange of protein complexes between the stacked and unstacked thylakoid regions. Therefore, to respond to varying environmental conditions that chloroplasts experience throughout a typical day, thylakoids should undergo some structural changes, at least at the scale of single protein complexes moving between membrane domains.


But how fluid are these thylakoid membrane domains? In other words, how mobile are the protein complexes and lipids within the plane of the membrane under short timescales? Do the components freely mix like a liquid, or are they more static like a gel? While plastoquinone must diffuse between PSII and cytochrome

*b<sub>6</sub>f*, it is unknown whether the larger photosynthetic complexes freely move around within their resident membrane domain or are fixed in space and experience only small oscillations (Kirchhoff 2014a). On the one hand, the low saturation of thylakoid lipids should facilitate the diffusion of protein complexes in the membrane; on the other, mobility might be restricted by molecular crowding from the high protein concentration (thylakoids are ~70% protein [Kirchhoff et al. 2002]) as well as electrostatic interactions between stacked thylakoids (Standfuss et al. 2005). Can current technology measure the degree of mobility for thylakoid protein complexes *in vivo*? And would changes in this mobility impact cellular physiology and photosynthesis? Does it matter whether thylakoid membranes are more fluid or more static?

Two electron microscopy (EM) techniques can directly visualize individual protein complexes within thylakoid membranes *in vivo*. Freeze-fracture EM has been used for decades to quantify the organization of photosynthetic complexes within fractured membrane planes (Ojakian and Satir 1974; Charuvi et al. 2016). More recently, cryo-electron tomography has extended this visualization to both sides of thylakoid membranes within the cell, providing a clear view of the lateral heterogeneity between membrane domains (Wietrzynski et al. 2020). However, these EM approaches can only provide frozen snapshots of protein complex organization, begging the question of how mobile these complexes are. Fluorescence recovery after photobleaching (FRAP) can provide information on protein mobility, and it has been used to determine that PSII complexes do not rapidly diffuse between separate grana stacks (Kirchhoff et al. 2008), which is consistent with lateral heterogeneity. However, the resolution of this technique is far too limited to follow the mobility of an individual PSII complexes within an appressed grana membrane. Spectroscopy can detect *in vivo* changes in the general ordering of thylakoid proteins, but it provides no spatial information on the movement of complexes. Only high-speed atomic force microscopy (AFM) combines the spatial resolution to map single complexes in a membrane with the temporal resolution to track their mobility (Clausen et al. 2014; Onoa et al. 2020). However, AFM is limited to measuring the topology of isolated membranes, which may not reproduce the thylakoid organization and fluidity found inside a living cell. Unfortunately, no single technique in use today can track the short-timescale mobility of individual protein complexes within native thylakoids.

Observations of PSII organization provide hints that this photosynthetic complex may experience changes in its mobility, and also underline the potential differences between isolated and native membranes. In cyanobacteria and red algae, PSII complexes bound along linear arrays of phycobilisomes must experience some restriction in their degrees of freedom. In isolated thylakoids of plants and diatoms, PSII has been observed in both a dispersed organization and arranged in two-dimensional pseudocrystalline lattices (Fig. 6) (Staehelin 1976; Dekker and Boekema 2005; Kirchhoff et al. 2007; Daum et al. 2010; Sznee et al. 2011; Levitan et al. 2019). These PSII lattices would further restrict the degrees of freedom, and they also interact with lattices in neighboring membranes to induce stacking of isolated thylakoids (Daum et al. 2010), limiting the mobility of PSII in two appressed membranes. *In vivo*, only the dispersed organization of PSII has been observed to date, with the exception of PSII lattices within dehydrated resurrection plants (*Craterostigma pumilum*) (Charuvi et al. 2015).

What would happen to PSII-related processes when the complexes are organized in a lattice within appressed thylakoids? How would the lattice affect plastoquinone diffusion to and



**Figure 6.** Does PSII organization and mobility change in vivo? Schematic illustrating the dispersed and pseudocrystalline lattice arrangements of PSII (blue), which could potentially modulate an inverse relationship between local PSII concentration and mobility. Whether such a transition in PSII organization is physiologically relevant in vivo remains to be explored.

from PSII? Are PSII able to leave the lattice when damaged? Are initiators (e.g. kinases; Depège et al. 2003) and effectors (e.g. proteases; Lindahl et al. 2000) located in the stroma able to reach them? Do PSII lattices receive less excitation from LHCII? If PSII lattices are physiologically relevant, perhaps a transition between mobile dispersed PSII and the static lattice configuration can help chloroplasts resist particularly harsh environmental conditions, such as dehydration of resurrection plants. The thylakoids may be locked in a nonphotosynthetic state that can activate again upon rehydration. Could such a change in thylakoid fluidity occur locally as a photoprotective mechanism to regulate excess excitation or to sequester subpopulations of damaged PSII? Due to the technical challenge of observing PSII dynamics in vivo, exploring these questions will require approaches that integrate structural observations, biophysical measurements, physiology, and molecular simulations.

## How does alternative electron flow contribute to the maintenance of robust photosynthesis?

(Written by Arthur R. Grossman)

Light absorption drives photosynthetic electron transport and the synthesis of reductant and ATP to energize  $\text{CO}_2$  fixation, growth, and other anabolic processes. However, photosynthetic antennae often absorb excess excitation energy, especially upon exposure of plants to high/fluctuating light intensities or stress (Saroussi et al. 2017). To cope with excess absorbed light energy, photosynthetic organisms have evolved mechanisms to dissipate this energy as heat through nonphotochemical quenching (NPQ), including energy-dependent quenching (qE), state transitions (qT), photoinhibition (qI), and zeaxanthin-dependent quenching that does not require a high  $\Delta\text{pH}$  across the thylakoid membranes (qZ) (Niyogi and Truong 2013).

Photosynthetic electron transport involves linear electron flow, which generates ATP and reducing power, and alternative electron flows (AEF), including cyclic electron flow and  $\text{H}_2\text{O}$ -to- $\text{H}_2\text{O}$  cycles in which  $\text{O}_2$  is reduced to  $\text{H}_2\text{O}$  (Fig. 7). Linear electron flow does not generate enough ATP to sustain photosynthesis, a deficit made up by the various AEF pathways. Additionally, AEF deposits protons in the thylakoid lumen that can help establish NPQ, impact the rate of linear electron flow, protect PSI from damage under high/fluctuating light (see the section below by Burlacot), and help balance the ATP:NADPH ratio, which can impact PSI donor and acceptor side regulation (Munekage et al. 2004; Allahverdiyeva et al.


2015a, 2015b). AEF reactions include (i) Mehler-type, (ii) NADPH:flavin oxidoreductase (FLV) catalyzed  $\text{O}_2$  reduction, designated pseudocyclic electron flow, (iii) plastoquinol terminal oxidoreductase (PTOX) activity, and (iv) chloroplast-to-mitochondria electron flow, which shuttles electrons between the organelles. The biochemical shuttles important for the flow of reductant between the chloroplast and mitochondria translocate oxaloacetate/malate (OMT; Fig. 7, #1a), malate/aspartate (Dao et al. 2022), and triose-P/Pi (TPT) (Fig. 7, #1b) (Huang et al. 2023), coupled with redox exchange reactions catalyzed by malate dehydrogenase (MDH) and glyceraldehyde 3-P dehydrogenase (GAPDH). We describe basic features of these pathways, with a focus on *Chlamydomonas*, and highlight some major unanswered questions about their role(s) in maintaining robust photosynthesis.

## Cyclic electron flow

Cyclic electron flow (Fig. 7, #2a, 2b) directs reducing equivalents from the PSI acceptor to the donor side, enabling proton deposition in the thylakoid lumen and the use of the proton gradient for ATP synthesis without generating reducing equivalents. Most photosynthetic organisms use two cyclic electron flow pathways, one involving the proton gradient generation 5 (PGR5) and the PGR-Like 1 (PGRL1) proteins (DalCorso et al. 2008), and the other the plastid Type I NDH (NADPH:flavin oxidoreductase) complex in vascular plants (Joet et al. 2001) or Type II NDH, designated NDA2, in green algae (Desplats et al. 2009). The Type I NDH has 11 chloroplast-encoded and >19 nucleus-encoded subunits and is similar to the bacterial/mitochondrial NADH:UQ respiratory complex (complex 1) (Peltier et al. 2016); it likely uses reduced ferredoxin (FDX) as the electron donor (Yamamoto et al. 2011). The PGR5/PGRL1 pathway also appears to involve PetO, FNR, and ANR2 (Buchert et al. 2018) that may function in a cyclic electron flow supercomplex (Joliot et al. 2022). Tobacco (*Nicotiana tabacum*) and *Arabidopsis* (*Arabidopsis thaliana*) with lesions in genes encoding chloroplast NDH subunits do not exhibit strong phenotypes under mild conditions (Shikanai et al. 1998). The *Chlamydomonas* Type II NDA2 transfers electrons from NAD(P)H to the PQ pool and becomes highly active during N deprivation (Saroussi et al. 2016).

## Mehler reaction

The Mehler reaction (Fig. 7, #3) involves noncatalyzed  $\text{O}_2$  reduction using PSI-derived electrons (Asada 1999). It can generate reactive oxygen species (ROS), including superoxides ( $\text{O}_2^-$ ) and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) that can cause damage to various molecules of the cell, or the  $\text{O}_2^-$  can be converted to  $\text{H}_2\text{O}$  by the



**Figure 7.** Diagram of photosynthetic electron transport in *Chlamydomonas*, two major pathways for exporting fixed carbon and/or reductant from chloroplasts, and alternative pathways for electron flows (AEF). Linear photosynthetic electron transport involves excitation of reaction centers (PSI, PSII) and extraction of electrons from H<sub>2</sub>O by the PSII O<sub>2</sub> evolving complex. Extracted electrons pass through PSII reaction centers, the plastoquinone (PQ) pool, Cytochrome b<sub>6</sub>/f (Cyt), plastocyanin (PC) and to PSI where they are used to generate reduced ferredoxin (FDX) and NADPH; the ATP synthesized by the ATP synthase (fueled by proton gradient across thylakoid membranes) and the NADPH are used to drive CO<sub>2</sub> fixation by the Calvin-Benson cycle. Major steps in these pathways are numbered in orange boxes. Fixed carbon and reductant are exported from chloroplasts by various shuttles; two major shuttles involve oxaloacetate/malate and triose-P/Pi exchange (OMT and TPT, respectively, 1a and 1b). Additionally, reducing electrons generated on the acceptor side of PSI or PSII can be routed through other AEF pathways (corresponding to 2 to 5) which include cyclic electron flow through both PGR5/PGRL1 (2a) and NDA2 (2b) pathways, the Mehler reaction (3) in which PSI-derived electrons are used to reduce O<sub>2</sub> and the ROS generated can be converted to H<sub>2</sub>O through superoxide dismutase and catalase/ascorbate peroxidase, pseudocyclic electron flow (4) in which electrons can be used to reduce O<sub>2</sub> to H<sub>2</sub>O through FLV flavodiiron proteins, and plastoquinol terminal oxidase (PTOX) catalyzed reduction of O<sub>2</sub> (5) associated with the acceptor side of PSII. Major pathways associated with AEF are depicted with orange arrows while black arrows indicate linear electron flow.

sequential activities of superoxide dismutase and ascorbate peroxidase/catalase. The Mehler reaction can also generate a  $\Delta$ pH across thylakoid membranes (Makino et al. 2002), although it is unclear how prevalent this reaction is in *Chlamydomonas*.

### Pseudocyclic electron flow

Pseudocyclic electron flow (Fig. 7, #4) involves electron transfer to PSI-associated flavodiiron proteins (FLVs), likely through FDX (Sétif et al. 2020), which are thought to directly reduce O<sub>2</sub> to H<sub>2</sub>O (Chaux et al. 2017). This reaction occurs in green algae, cyanobacteria, mosses, liverworts, and gymnosperms, but not in angiosperms, and is especially important immediately after dark acclimation when the Calvin-Benson cycle is not activated, and under fluctuating light conditions (Allahverdiyeva et al. 2013). A more detailed discussion of the flavodiiron proteins is given in the section below by Allahverdiyeva et al.

### Plastoquinol terminal oxidoreductase

PTOX, associated with chlororespiration (Fig. 7, #5), uses electrons from the PQ pool to reduce O<sub>2</sub> to H<sub>2</sub>O (Peltier et al. 2010). These electrons can come directly from PSII or from NADPH to the PQ pool via the NAD(P)H-PQ reductase; Type I NDH in plants and NDA2 in green algae (e.g. *Chlamydomonas*). Potential roles of PTOX include protection of cells during fluctuating light (Nawrocki et al. 2019) and PQ pool over-reduction, which can elicit

PSII charge recombination and the generation of singlet O<sub>2</sub>. *Chlamydomonas* has two PTOX isoforms (PTOX1, PTOX2) (Houille-Vernes et al. 2011), with PTOX1 likely involved in regenerating oxidized PQ for phytoene desaturation (Foudree et al. 2012).

### Chloroplast-to-mitochondria electron flow

The transfer of reducing equivalents between chloroplasts and mitochondria, here referred to as chloroplast-to-mitochondria electron flow (Fig. 7, #1a,1b), is also integral for controlling cellular energetics and redox conditions in the light, and for sustaining dark anabolic processes (Dang et al. 2014). Chloroplast-mitochondria interactions were suggested in studies using inhibitors of mitochondrial electron flow and mutants in chloroplast ATP synthase (Lemaire et al. 1988) and respiration (Cardol et al. 2009). The integration between chloroplast and mitochondrial electron flow (He et al. 2023) helps coordinate the two pathways, balancing the production of reductant and ATP, regulating the metabolism of phosphate, nitrogen, and carbon compounds, and potentially managing accumulation of ROS to help sustain chloroplast electron transport and CO<sub>2</sub> fixation. Major translocators for chloroplast-to-mitochondria electron flow have been proposed to involve OMTs (malate/oxaloacetate, malate/aspartate) and TPTs. Plants and algae have multiple OMTs and TPTs, which are bidirectional transporters that function based on mass action and can move carbon and reductant out of chloroplasts (as these

metabolites accumulate in the stroma), delivering them to other cellular compartments, including mitochondria. Reductant delivered by these transporters to mitochondria can serve as substrate for respiratory electron transport and reduction of O<sub>2</sub> through cytochrome oxidase or alternative oxidases (AOX), which are present in both plant and algal mitochondria. Inhibitor studies (myxothiazol, an inhibitor of the mitochondrial cytochrome *bc*<sub>1</sub> complex, and salicylhydroxamic acid, an inhibitor of the mitochondrial alternative oxidase) also suggest that the AOX pathway may generate some chemical bond energy/ATP that can support photosynthetic CO<sub>2</sub> fixation through the formation of electrochemical potential (although AOX does not pump protons), but not as much as the cytochrome oxidase pathway (Peltier et al. 2023). The AOX pathway is cyanide insensitive, enables continued operation of glycolysis and the TCA cycle under stress conditions when there is restricted electron flow through cytochrome oxidase, balances C:N, ATP:ADP, NAD(P)H:ATP ratios, and limits hyper-reduction of the respiratory chain and accumulation of ROS (Rogov et al. 2014; Kaye et al. 2019).

While OMTs are thought to mainly shuttle reducing equivalents between chloroplasts and other cellular compartments, TPTs may mainly export fixed carbon synthesized in chloroplasts (Huang et al. 2023). Chloroplast TPTs can transport triose-P [glyceraldehyde 3-P (GAP), dihydroxyacetone-P (DHAP), and 3-phosphoglycerate (3-PGA)] in a counter exchange for cytosolic inorganic phosphate (Pi), which resupplies the chloroplast with Pi, allowing for continued photosynthetic electron flow and CO<sub>2</sub> fixation. TPTs are part of the pPT plastid translocator family that can serve as antiporters of phosphorylated C<sub>3</sub>, C<sub>5</sub>, or C<sub>6</sub> compounds with Pi (Flügge et al. 2003; Bockwoldt et al. 2019). In plants, trioses exported from chloroplasts are used to synthesize sucrose and other metabolites and fuel respiratory activity. Plants also harbor other pPT subfamilies, including those for glucose 6-P (GPTs) (Kammerer et al. 1998), xylulose-P (pentose-P) (Eicks et al. 2002), and phosphoenolpyruvate (PEP) (PPTs) (Fischer et al. 1997); PEP can be imported into C<sub>3</sub> plant plastids and used as substrate to fuel fatty acid biosynthesis or the shikimate pathway, and exported from C<sub>4</sub> plant plastids (Streatfield et al. 1999; Prabhakar et al. 2010). In some plants, diminished TPT levels do not yield a strong phenotype because the loss of plastid TPT activity can be compensated for by accumulation of a transitory starch pool that can be rapidly degraded (Hausler et al. 1998, Walters et al. 2004) to products (e.g. hexose-P) exported from chloroplasts and used in other cellular compartments. *Chlamydomonas* TPT3 was shown to likely be most active in routing triose-P out of chloroplasts; *tpt3* null mutants exhibited aberrant photosynthesis and highly reducing conditions in chloroplasts (Huang et al. 2023).

### Contributions of AEF pathways

The relative importance of the different AEF pathways in generating the ATP needed to sustain CO<sub>2</sub> fixation has been evaluated using electron transport inhibitors and mutants. Cyclic electron flow, pseudocyclic electron flow, and chloroplast-to-mitochondria electron flow activities were quantified in *Chlamydomonas* and shown to have compensatory activities, with each able to provide a large fraction of the energy needed to sustain photosynthesis (Peltier et al. 2023); the most energetically efficient pathway was chloroplast-to-mitochondria electron flow. However, using mutants and pathway inhibitors to quantify individual pathway activities may not accurately reflect *in vivo* wild-type activities since a loss of any of these pathways *in vivo* may alter the contributions

of the remaining pathways, and in some cases the inhibitors may not be completely penetrant.

### Questions concerning integration of AEF activities

Photosynthetic electron flow and energy transfer are highly complex processes (Fig. 7) and there is ongoing debate about the importance of the different AEF pathways. Electron flow and energy transfer through these pathways largely occur through metabolic flux driven by mass action. This means that blocking one “road” can increase flux through another. Organisms can make adjustments by modifying gene expression or other regulatory phenomena. Some of the unknown questions include the following.


1. Can we measure the precise contributions of each AEF pathway *in vivo* without disrupting (mutants, inhibitors) the system?
2. What conditions govern the utilization of specific AEF activities and the proportion of each?
3. Which components of AEF complexes (e.g. cyclic electron flow) are involved in catalytic activity and regulation?
4. How do subunits of individual AEF complexes interact; what controls these interactions?
5. What components of AEF pathways are post-translationally modified; how do such modifications impact protein structure, interactions, and activities?
6. Why are there two different NDH pathways (Type I and II); do differences in these pathways reflect their ability to use different substrates?
7. What are the functions of the chloroplast OMTs and TPTs in driving chloroplast-to-mitochondria electron flow and what regulates their activities?
8. How are the activities of transporters of phosphorylated sugars on the membranes of various cellular compartments integrated with chloroplast sugar-P transport?

With an initial plunge into the waters of AEF and associated metabolite trafficking, we are just beginning to realize the depth of those waters.

### Do flavodiiron proteins have multiple physiological roles in photosynthetic organisms?

(Written by Yagut Allahverdiyeva, Henna Mustila, and Lauri Nikkanen)

Flavodiiron proteins (FLVs) play a crucial role in some photosynthetic organisms as an alternative electron transport pathway, allowing growth in fluctuating light intensities by providing a valve for the safe dissipation of excess electrons in the photosynthetic electron transport chain. FLVs drive a four-electron reaction reducing O<sub>2</sub> to water downstream of PSI. This reaction is known as the Mehler-like reaction due to constituting a water–water cycle similarly to the Mehler reaction, or pseudocyclic electron transport (Fig. 7 #4, and see above section by Grossman). FLVs are found in various photosynthetic organisms ranging from cyanobacteria to vascular plants, but are absent in red algae and angiosperms. Cyanobacterial Flv1 and Flv3 have structural and functional similarities to FlvA and FlvB in algae and plants (Allahverdiyeva et al. 2015a, 2015b, Ilik et al. 2017). The reason for the loss of FLVs from angiosperms during evolution remains elusive. However, it may



**Figure 8.** Function and regulation of FLVs in *Synechocystis* and *Anabaena*. **A)** In *Synechocystis*, Flv1/Flv3 and Flv2/Flv4 hetero-oligomers function in light-driven  $O_2$  photoreduction and their expression level is dependent on carbon availability. In addition, Flv3/Flv3 homo-oligomer catalyzes a distinct, unspecified reaction than the  $O_2$  photoreduction. **B)** The reversible association of different Flv1/3 and Flv2/4 hetero-oligomers with the thylakoid membrane of *Synechocystis* is controlled by the extent of the trans-thylakoid proton gradient. The + symbols refer to positively charged protons. See text for details. **C)** In filamentous *Anabaena*, Flv1A/Flv3A function in vegetative cells and Flv3B/Flv3B in heterocysts in  $O_2$  photoreduction, the latter configuration maintaining microoxic conditions to protect the  $N_2$ -fixing enzyme nitrogenase in the light. The roles of Flv2/Flv4 and Flv1B/Flv1B in *Anabaena* are yet to be elucidated.

relate to increased efficiency of other electron sinks and protective mechanisms such as nonphotochemical quenching, reactive oxygen species scavenging systems, and cyclic electron transport, which would have made retention of energy-wasting FLVs redundant or even disadvantageous (for discussion on this, see e.g. Alboresi et al. 2019b).

In the reference cyanobacterium *Synechocystis* (*Synechocystis* sp. PCC6803), the Flv1/Flv3 hetero-oligomer catalyzes strong but transient  $O_2$  photoreduction in the first minute of illumination or during increased light intensity under ambient  $CO_2$  (Santana-Sánchez et al. 2019), providing essential electron sink capacity during the initial phase of photosynthetic induction before full activation of the Calvin–Benson cycle (Fig. 8A). Accordingly, *flv* mutants (deficient in Flv1 or Flv3 or homologs FlvA and FlvB) demonstrate strong growth retardation under fluctuating light, where low background light is repeatedly interrupted by high-light pulses, causing strong acceptor-side limitation of PSI (Allahverdiyeva et al. 2013; Gerotto et al. 2016; Chaux et al. 2017; Jokel et al. 2018). *Synechocystis* Flv mutants show a less pronounced phenotype under milder fluctuating light conditions (higher background light) at pH 8.2, and no growth phenotype was observed at pH 7.5 (Mustila et al. 2016), suggesting an external pH-dependent effect on FLVs and growth that remains poorly understood.

In addition to Flv1 and Flv3, some  $\beta$ -cyanobacteria possess Flv2 and Flv4 proteins (Helman et al. 2003). These proteins also participate in Mehler-like reactions downstream of PSI, but with different kinetics, catalyzing  $O_2$  photoreduction that persists beyond the initial phases of photosynthetic induction (Santana-Sánchez et al. 2019; Fig. 8A). Unlike Flv1 and Flv3, Flv2 and Flv4 are not crucial during sudden increases in light intensity. In *Synechocystis*, Flv2 and Flv4 show strong downregulation under low light, high  $CO_2$  and high pH conditions (Eisenhut et al. 2012; Santana-Sánchez et al. 2019). Deletion of Flv2 and Flv4 alters the phenotype under high light, and overexpression of *flv4-2* operon provides resistance to high light (Bersanini et al. 2014). Notably, the presence of these proteins within an operon alongside Sll0218 suggests that some phenotypic changes could be attributed to the role of this small protein in PSII assembly and stabilization (Bersanini et al. 2017).

Heterocyst-forming  $N_2$ -fixing cyanobacteria, such as *Anabaena* (*Anabaena* sp. PCC 7120), possess six FLVs (Fig. 8C). Flv1A and Flv3A are specific to vegetative cells and are crucial for survival under FL conditions. Interestingly, deletion of Flv3A affects  $N_2$  fixation and  $H_2$  metabolism in heterocysts, specialized  $N_2$  fixing cells, via downregulation of uptake hydrogenase, enabling  $H_2$  production even under oxic conditions (Santana-Sánchez et al. 2023).

Investigations into the roles of Flv2 and Flv4 in *Anabaena* are yet to be conducted, although their expression pattern under different conditions suggests a function similar to that in *Synechocystis*. Flv1B and Flv3B are specific to heterocysts (Ermakova et al. 2013, 2014). Unlike other cyanobacterial FLVs, Flv3B can function as a homo-oligomer in the Mehler-like reaction, maintaining microoxic conditions under light. Importantly, while Flv1B is not involved in the Mehler-like reaction, its deletion strongly impacts  $N_2$  fixation, indicating a role in  $N_2$  metabolism (Ermakova et al. 2014).

### What are the functional oligomeric states of FLVs?

All FLVs contain an N-terminal metallo- $\beta$ -lactamase-like domain harboring an  $O_2$  and/or NO reducing nonheme diiron center and a C-terminal flavodoxin-like (FMN) domain (Vicente et al. 2008). Oxygenic photosynthetic organisms contain an additional NAD(P)H:flavin reductase-like domain fused in the C-terminus. *In vitro* studies and modeling suggested that FLVs function in a head-to-tail homo-oligomeric configuration (Vicente et al. 2008) enabling inter-subunit electron transfer, between Fe-Fe of one monomer and FMN of the other. In *Synechocystis*, an *in vivo* hetero-oligomeric configuration (Flv1/3 and Flv2/4) is important for  $O_2$  photoreduction: *Synechocystis* strains lacking Flv1 but overexpressing Flv3 (thus Flv3/3 configuration) or lacking Flv3 but overexpressing Flv1 (Flv1/1) did not demonstrate  $O_2$  photoreduction (Mustila et al. 2016). However, having overexpressed homo-oligomers of Flv3 or Flv1 partly rescued the phenotype under FL, suggesting that homo-oligomers can function in unknown metabolic pathways and play a role in photoprotection that is independent of  $O_2$  photoreduction. Moreover, in *Anabaena*, the deletion of Flv1A did not result in total impairment of the Mehler-like reaction in air-level  $CO_2$  conditions where Flv2 and Flv4 are expressed (Santana-Sánchez et al. 2023). This suggests that Flv3A can catalyze  $O_2$  photoreduction in an Flv2/4-dependent manner either as a homo-oligomer interacting with Flv2/4 or by forming oligomers with Flv2 or Flv4.

### Can FLVs in photosynthetic organisms catalyze NO reduction?

FLVs can reduce NO into  $N_2O$  in anaerobic bacteria (Vicente et al. 2008) and in *Chlamydomonas reinhardtii* (Burlacot et al. 2020). We cannot exclude a similar function in cyanobacteria, although the efficiency of this reaction as a sink is disputable. However, it may affect cell metabolism via signaling, and could be aligned with alternative electron donors. Recent data show that *Synechocystis* Fed1, the most abundant of 11 ferredoxins in this cyanobacterium, can interact with and act as a donor to the Mehler-like reaction catalyzed by Flv1/3 and Flv2/4 (Nikkanen et al. 2020, 2023; Sétif et al. 2020). However, *in vitro* studies demonstrated that NAD(P)H is a donor for recombinant Flv1, Flv3, or Flv4 homo-oligomers (Vicente et al. 2008; Shimakawa et al. 2015; Brown et al. 2019), albeit with relatively low electron transfer rates. Interestingly, a protein-protein interaction was detected between Flv3 (but not Flv1) and the large isoform of FNR ( $FNR_L$ ) (Nikkanen et al. 2023). Flv3/ $FNR_L$  interactions could occur to facilitate electron transfer from NADPH to Flv3 homo-oligomers, functioning in a pathway to reduce an unknown acceptor, such as NO. Flv1-independent activity of Flv3 is also supported by 6-fold higher protein amount of Flv3 in comparison to Flv1 (Jackson et al. 2023). However, the physiological relevance of these interactions and Flv3-homo-oligomer-dependent electron transfer pathways remain unresolved.

### How is FLV activity regulated?


The Mehler-like reaction displays distinct kinetics where Flv1/3-dependent  $O_2$  photoreduction is transiently induced and then inactivated during the first minute of illumination. This strongly suggests that FLV activity is subject to post-translational regulation (Alboresi et al. 2019a). Recently, we proposed a model for pH-dependent regulation of FLV hetero-oligomer activity (Nikkanen et al. 2023), where alkalization of the cytosol during photosynthetic induction results in Flv1/3 hetero-oligomers being repelled from the thylakoid membrane due to their surface charge becoming more negative and their activity being thus decreased (Fig. 8B). Flv2/4 remains associated with the thylakoids longer after photosynthetic induction (i.e. in a more alkaline cytosol) due to large positively charged patches in Flv4. While this model explains many features of the observed kinetics of FLV activity, FLVs are likely subject to additional layers of post-translational regulation. The fact that we see a transient reduction of ferredoxin at all during dark-to-light transitions suggests that Flv1/3 hetero-oligomers are not yet fully primed in darkness, despite cytosolic pH being neutral and FLV hetero-oligomers being associated with the thylakoids. Therefore, it is likely that an additional unknown signal is required for the full light-dependent activation of Flv1/3 hetero-oligomers. Indeed, FLVs have been suggested to be redox-regulated via conserved cysteine residues, some of which in Flv1 and Flv3 are light-dependently reduced (Guo et al. 2014). It remains to be elucidated if thiol modulation or other post-translational modifications, such as phosphorylation, play a role in regulating FLV activity.

An additional question concerns how FLV activity is regulated in low versus high  $CO_2$  conditions. In *Synechocystis*, in high  $CO_2$ , although Flv1 and Flv3 are downregulated, and Flv2 and Flv4 are not expressed (Santana-Sánchez et al. 2019), FLV-dependent  $O_2$  photoreduction activity is sustained at a higher level than in low  $CO_2$ . This suggests a regulatory mechanism that connects FLV activity to carbon metabolism and possibly to carbon concentrating mechanisms.

### How is photosynthesis regulated during light fluctuations?

(Written by Adrien Burlacot)

Photochemical reactions happen extremely fast (on the picosecond timescale), but the redox reactions involved in the transport of electrons are orders of magnitude slower, and the metabolic reactions of carbon metabolism are even slower. This timescale discrepancy makes meeting metabolic energy requirements with light-energy-driven PS activity challenging to attain when light intensity is variable. In extreme cases of light fluctuation, photosynthetic electron generation can overflow electron transport or metabolic capacity, leading to PS degradation (Sejima et al. 2014) and potentially cell death. In many natural contexts, light rapidly fluctuates between a sub-saturating (low light, LL) to an over-saturating level (high light, HL) (Fig. 9A), and cells must cope with either excessive or insufficient amounts of energy and quickly switch between those states as light changes. The electron transport chain is a central hub of regulation in this context. Despite decades of investigations, how photosynthesis is dynamically controlled to provide chemical energy without becoming harmful during light fluctuations remains a thriving area of research (Morales and Kaiser 2020). This question has recently regained traction with groups showing that tuning the dynamics of light energy management can improve the productivity of



**Figure 9.** Acclimation mechanisms to light fluctuations. **A)** Light fluctuations generated by diurnal cycle and cloud coverage on top of a field canopy in June 2021. Data shown are light intensity of the photosynthetically active radiations throughout the time of the day from the Surface radiation Budget Network (SurfRAD, NOAA) at the Bondville Station, Illinois (40°N latitude; 88°W longitude). **B)** Main molecular mechanisms involved in tuning photosynthetic electron transport during artificial light fluctuations. The luminal pH controlled by ion transport (mediated by KEA3, VCCN1, and the ATPase) and alternative electron transport (mediated by PGR5/L1 and FLVs), activates NPQ mechanisms and controls the photosynthetic electron flow at the cytochrome  $b_6f$  (Cyt  $b_6f$ ) level. Phosphorylation of LHCII, which dynamically relocates LHCII to PSI, and redox exchange between the chloroplast and the rest of the cell are also involved in the dynamic regulation of photosynthetic electron transport. Shown in blue arrows are photosynthetic electron transport pathways. **C)** Example of characteristics of light fluctuations that can be explored to define the domain of importance of molecular mechanisms involved. The definition of such domains might help in predicting the set of mechanisms relevant to survive in a more complex light pattern.

photosynthetic organisms (Kromdijk et al. 2016; De Souza et al. 2022; Perin et al. 2023). Previous sections provided an overview of alternative electron flow pathways (Grossman) and the role of the pathway mediated by FLVs in some organisms (Allahverdiyeva et al.). Here, I discuss questions related to how organisms regulate the dynamics of which pathways and mechanisms are employed under fluctuating light conditions in different environments.

Many dynamic regulatory mechanisms of photosynthetic electron generation and management are triggered by low luminal pH, which modulates electron transport at the level of the cytochrome  $b_6f$  (Malone et al. 2021) and activates nonphotochemical quenching (NPQ) of excess absorbed light energy. Since the discovery of the role of thylakoid protein phosphorylation in

response to 5 min LL/1 min HL treatment (Fig. 9B) (Tikkanen et al. 2010), similar artificial light patterns have been extensively used to identify proteins involved in the adjustment of the energetic status of the electron transport chain when light fluctuates. In model species of angiosperms, mosses, microalgae, and cyanobacteria (Fig. 9B) those include: (i) proton sensing proteins involved in NPQ which limit photochemistry and protect PS during HL (Cantrell and Peers 2017; Roach 2020; Steen et al. 2022; Niu et al. 2023); (ii) ion exchange across the thylakoid membrane mediated by VCCN1 and KEA3, which balances the proton motive force and controls NPQ and electron flow during the few minutes following a light change (Armbruster et al. 2016; Herdean et al. 2016); (iii) the plastid terminal oxidase (PTOX), which contributes to the redox poise of the thylakoid during the LL/dark phase of

fluctuation (Nawrocki et al. 2019), (iv) cyclic electron flow controlled by PGR5 and PGRL1 proteins which tunes lumenal pH and electron flow during the first minutes of HL (Yamamoto et al. 2016; Yamamoto and Shikanai 2018; Storti et al. 2019), (v) FLVs, which dissipate the overflow of electrons and contribute to the low lumenal pH upon HL exposure within a few seconds by reducing O<sub>2</sub> to water (Allahverdiyeva et al. 2013; Gerotto et al. 2016; Chaux et al. 2017; Jokel et al. 2018), and (vi) redox shuttles which rebalance the redox poise between the organelles (Yokochi et al. 2021).

Despite this extensive amount of knowledge, the complete set of proteins involved during light fluctuations is likely just beginning to be revealed. Indeed, acclimation mechanisms to HL or LL involve (i) posttranslational regulation of protein activity (Allorent et al. 2013) (on a timescale of milliseconds to tens of minutes), (ii) translation of mRNA (Durnford et al. 2003) (on a timescale of minutes), (iii) changes of mRNA levels (Mettler et al. 2014), (iv) structural changes of organellar membranes (Kirchhoff 2014b) or (v) leaf orientation or morphology (Feng et al. 2019) (on a timescale of minutes to hours/days). Hence, the preceding time in LL or in HL, the intensities of LL and HL, the shape of the light transient as well as how frequent light is shifting likely have a major impact on the identity and role of mechanisms involved in acclimation. The use of just one light fluctuation regime becomes thus limited both for discovering new proteins involved and for identifying domains where known proteins are important.

How do different light fluctuation characteristics change which mechanisms are involved in photosynthesis regulation? Reductionist approaches hold great promises in that regard. One of them, introduced by Nedbal and Březina (2002) and recently used on mutated strains by Shimakawa and Miyake (2018); Steen et al. (2022), and Niu et al. (2023) involves assessing the response of mutants impaired in photosynthetic regulatory processes to artificial periodic illumination (square-like or sine-like). Varying the periodicities used over short-term episodes of light fluctuations allows us to define the periodicity domain of photosynthetic regulatory processes, which are periodicity limits below or above which a regulatory mechanism cannot dynamically respond to a periodic fluctuation. Defining the periodicity domain of an acclimatory mechanism might allow predicting its importance during a more complex light fluctuation harboring a wide set of fluctuation periodicities. Note here that while this method is common in engineering sciences and relates to frequency-domain analysis, I refer here to the periodicity domain for notation simplicity. This method has allowed the unraveling of periodicity domains of proteins involved in NPQ: (i) in the green algae *Chlamydomonas*, NPQ magnitude and dynamics depend more on the light-harvesting complex stress-related 3 protein (LHCSR3) for a 1 min periodicity than for a 10 min periodicity (Steen et al. 2022), and (ii) in the angiosperm *Arabidopsis*, NPQ does not respond to dynamic light changes below a periodicity of 10 s while for periodicities of 10 to 60 s, the protein PsbS is critical for NPQ response; for longer periodicities, both PsbS and changes in the xanthophyll cycle contribute to the NPQ response (Niu et al. 2023). However, while defining periodicity limits of acclimatory processes can prove important, it remains limited to short-term episodes of light fluctuation and misses long-term acclimatory mechanisms. Another approach consists of growing/acclimating the photosynthetic organism under different patterns of light fluctuation and measuring how photosynthetic parameters are affected by the growth/acclimation conditions. Such an approach can reveal emergent phenomena in new mutants

(Peng et al. 2020) and has recently revealed that (i) the average growth light intensity affects the quantum efficiency of PSII and the dynamics of ion transport across the thylakoids upon HL while, (ii) the variability of growth light changes the levels and de-epoxidation state of the xanthophyll-cycle pigments violaxanthin, antheraxanthin, and zeaxanthin (von Bismarck et al. 2023). Future work combining those two approaches with other key characteristics of light fluctuations will be important to understand how the variability of light influences the mechanisms involved in acclimation to light.

What happens in a natural light environment? While fluctuations of light intensity in nature can be complex (Fig. 9A), reductionist methods like the ones discussed above are powerful and should allow, when used systematically, the prediction of the identity and activity of acclimatory processes involved depending on key characteristics of light fluctuations (Fig. 9C). The periodicity of light fluctuation is a clear characteristic of natural relevance, as under a forest canopy, the movement of leaves forms gaps that let high-intensity sunlight pass typically during less than 1.6s (Percy 1990) while, at the top of the canopy, HL exposure typically lasts for a few minutes (Smith and Berry 2013). Such periodicity variability might even be larger in aquatic environments where rippling caustic effects of the water surface fluctuate light with periodicities of less than 100 ms (Schubert et al. 2001). While other light fluctuation characteristics influencing light acclimation mechanisms like average, maximal, and minimal light intensity have been explored to some extent (Allahverdiyeva et al. 2013; Jokel et al. 2018; von Bismarck et al. 2023); identifying a subset of key light characteristics that describes natural light fluctuations in different environment and quantifying their range remains to be done. An interesting possibility here is to apply signal decomposition methods currently used to study the output of photovoltaic panels in electrical engineering (Meyers and Boyd 2023) to define what are the most relevant light intensity characteristics present in different natural light environments.

Importantly, the presence or the absence of proteins regulating photosynthesis in different species can vary widely and this diversity and its implication are critical to study. For example, while FLVs are required for acclimation to 1min HL/5min LL fluctuations in many phototrophs, they are absent in angiosperms (Yamamoto et al. 2016), the latter most likely relying on other mechanisms to cope with sudden HL transients. The development of knowledge on the diversity of mechanisms across species (Sello et al. 2019), together with modeling the potential interaction of different mechanisms (Li et al. 2021a, 2021b), might allow molecular breeding of plants for adaptation to their local light environment in conditions where acclimation to light limits growth (Perin et al. 2019).

How is photosynthesis regulated during light fluctuations? The development of new tools (Cruz et al. 2016) and methods (Nedbal and Lazár 2021) to systematically look at different aspects of this question will bring new perspectives to the complex regulations of photosynthesis under light fluctuations. And the answers will most likely depend on which characteristics of light fluctuation are investigated.

## How can photorespiration impact stomatal opening?

(Written by Michael Hodges)

Photosynthetic CO<sub>2</sub> assimilation in embryophytes requires the uptake of atmospheric CO<sub>2</sub> into the leaf. This is achieved by stomata, specialized epidermal structures composed of two guard


cells. They possess partially overlapping signal transduction networks controlling stomatal pore size by modulating guard cell turgor pressure in response to abiotic and biotic cues including light, CO<sub>2</sub>, water potential, temperature, ozone, and pathogens. In this way, plants balance CO<sub>2</sub> uptake and water loss with respect to fluctuating environmental conditions. Stomata are a target to be considered when designing future-proofed crops (Nguyen et al. 2023), therefore a complete understanding of how stomatal opening and closure are regulated is important. Although there is abundant knowledge concerning guard cell signaling networks and the H<sup>+</sup>-pumps, H<sup>+</sup>-coupled ion transporters, and ion channels involved in controlling stomatal movements (Saito and Uozumi 2019), how plant metabolism plays a role is less clear (Lemonnier and Lawson 2023). Mitochondrial activity (Nunes-Nesi et al. 2007) and cytosolic ATP (Lim et al. 2022) appear to be important for stomatal opening. Low blue light perceived by PHOT1/2 photoreceptors is involved in early morning guard cell starch degradation required for stomatal opening (Horrer et al. 2016). Photosynthesis has been shown to promote stomatal opening but its relative importance and cellular location (mesophyll cells versus guard cells) are still open to debate (Lemonnier and Lawson 2023). Recently, a role for photorespiration in stomatal functioning was suggested from experiments using photorespiratory mutants (Eisenhut et al. 2017; Flügel et al. 2017; Duminil 2019). If this is so, then how do photorespiratory cycle activity and/or photorespiratory-associated metabolites interfere with processes and signaling pathways that control stomatal opening/closure?

So what is photorespiration and which photorespiratory metabolites have the potential to influence stomatal opening/closure? Photorespiration is an essential metabolic process taking place in the light. It begins in chloroplasts with the oxygenase activity of Rubisco producing 2-phosphoglycolate (2-PG), which can inhibit the Calvin–Benson–Bassham cycle enzymes triose-phosphate isomerase and sedoheptulose 1,7-bisphosphate

phosphatase as well as glycolytic phosphofructokinase (Flügel et al. 2017 and references therein). The photorespiratory cycle spans four cell compartments (chloroplasts, peroxisomes, mitochondria, cytosol) and requires eight enzymes to transform toxic 2-PG into useful 3-phosphoglycerate. To accomplish this, photorespiration leads to the production of H<sub>2</sub>O<sub>2</sub> and NADH while liberating CO<sub>2</sub> and ammonia during serine formation from glycine in mitochondria (reviewed in Eisenhut et al. 2019).

As early as the 1960s, short-term pharmacological studies using photorespiratory enzyme inhibitors showed a correlation between photorespiration and stomatal opening in the light. The addition of hydroxysulfonate (an inhibitor of glycolate oxidase, a peroxisomal enzyme producing H<sub>2</sub>O<sub>2</sub> and glyoxylate from glycolate) to tobacco leaf disks (Zelitch and Walker 1964) and aminoacetonitrile (an inhibitor of glycine decarboxylase, a mitochondrial complex producing CO<sub>2</sub>) to rice leaves (Cho et al. 1987) provoked stomatal closure in the light. An association between photorespiration and stomatal opening/closure was further suggested because the inhibitory effect of aminoacetonitrile did not occur in detached rice leaves under low photorespiratory O<sub>2</sub> air conditions and aminoacetonitrile application to maize (*Zea mays*) leaves (a C<sub>4</sub> plant with low photorespiration) did not alter transpiration (Cho et al. 1987).

A renewed interest in photorespiration and stomata opening/closure has come from recent observations made using Arabidopsis photorespiration mutants (Fig. 10). Compared to wild-type (WT) plants, mutants lacking either phosphoglycolate phosphatase (*pglp1*), or serine hydroxymethyltransferase (*shm1*), or glycylate kinase (*glyk1*) showed higher and lower transpiration rates at high CO<sub>2</sub> (10,000 ppm) and in ambient air, respectively (Eisenhut et al. 2017). This was later partially confirmed by direct stomatal aperture measurements using epidermal peels of *pglp1* leaves and light microscopy (Duminil 2019). Arabidopsis PGLP1 antisense repression also led to lower stomatal conductance (*gs*) and transpiration rates when plants were transferred from high



**Figure 10.** Potential connections between photorespiration and metabolic processes relevant to stomatal opening/closure. In the light, Rubisco is involved in both photosynthetic CO<sub>2</sub> and photorespiratory O<sub>2</sub> assimilation. Photorespiration produces a number of metabolites that have the potential to impact stomatal opening and closure. 2-PG can reduce starch biosynthesis and therefore blue light/PHOT-induced malate from starch degradation and thus limit early morning stomatal opening. 2-PG might also inhibit glycolysis and malate biosynthesis required for stomatal opening while photorespiratory CO<sub>2</sub> and 3-phosphoglycerate (3-PGA) could be used to produce malate via PEP carboxylase (PEPC) activity for stomatal opening. Photorespiration also produces H<sub>2</sub>O<sub>2</sub> and serine, and both could be linked to stomatal closure either as a signaling molecule or via the production of O-acetylserine required for sulfate-induced ABA biosynthesis, respectively. Photorespiration also negatively impacts photosynthetic activity thus reducing the capacity to produce starch, sugars, and ATP that are important for stomatal opening.

to ambient CO<sub>2</sub> (Flügel et al. 2017). On the other hand, PGLP1 overexpression gave higher *g<sub>s</sub>* values and transpiration rates in 40% O<sub>2</sub> air; a high-photorespiration condition that increased leaf 2-PG amounts (Flügel et al. 2017). Lines overexpressing PGLP1 also maintained high *g<sub>s</sub>* levels at an elevated temperature and after drought stress, conditions associated with increased photorespiration (Timm et al. 2019). Greenhouse-grown tobacco overexpressing glycine decarboxylase (GDC)-H subunit also exhibited higher *g<sub>s</sub>* although this was not observed in the field (López-Calcano et al. 2019). By analyzing epidermal peels of complemented Arabidopsis *sh1* lines, SHM activity was found to impact salt- and ABA-induced stomatal closure (Liu et al. 2019).

So how can photorespiration impact stomatal opening/closing? There are a number of possible mechanisms that require investigation (Fig. 10).

- Transcriptional regulation could be involved since several photorespiratory mutants have deregulated genes involved in stomatal closure including *ABSCISIC ACID INSENSITIVE 1/2* (*ABI1/2*) (ABA biosynthesis), and *RESPIRATORY BURST OXIDASE HOMOLOGS D/F* (*RBOHD/F*) (reactive-oxygen production) and blue-light-induced stomatal opening (*PHOT1/2*) (Eisenhut et al. 2017). The signals and mechanisms involved are not yet known.
- Photorespiration could act by indirectly or directly impacting photosynthetic electron transfer and as a consequence ATP biosynthesis, which is required for stomatal opening. Photosynthetic processes have been shown to be affected by reduced photorespiratory cycle activity in mutants, including an attenuation of photosynthetic CO<sub>2</sub> assimilation, modification of PSII redox potential, and an increase in energy dissipation mechanisms (Timm et al. 2012; Dellero et al. 2015, 2016; Messant et al. 2018). Indeed, a correlation between *g<sub>s</sub>* and photosynthesis exists (Nunes-Nesi et al. 2007; Lu et al. 2014).
- Changes in photorespiratory activity could also provoke metabolic regulation by interacting with and perhaps modulating metabolic pathways relevant to stomatal movements including the biosynthesis of starch, malate, and ABA.
- Photorespiration could impact blue-light-induced stomatal opening at the beginning of the day. It has been proposed that enzyme inhibition by 2-PG is a control loop adjusting stomatal conductance to photosynthesis-related C-fluxes by altering C-allocation between RuBP regeneration and starch synthesis (Flügel et al. 2017). A correlation between PGLP activities, 2-PG levels, *g<sub>s</sub>*, and transpiration rates in PGLP transgenic lines (Eisenhut et al. 2017; Flügel et al. 2017; Timm et al. 2019) and reduced end-of-day leaf starch levels in photorespiratory mutants (Timm et al. 2012) support this hypothesis.
- Guard cell osmoregulation requires starch breakdown, triose-phosphate export from chloroplasts, and cytosolic PEP carboxylase (PEPC) activity for glycolytic malate production (Gehlen et al. 1996). Photorespiration produces 3-PGA and liberates CO<sub>2</sub> and therefore this could fuel glycolysis and PEPC activity to promote guard cell malate biosynthesis and stomatal opening. On the other hand, a build-up of photorespiratory 2-PG might inhibit glycolysis and favor stomatal closure.
- Sulfate-induced stomatal closure upon drought is associated with guard cell-specific ABA biosynthesis (Batoool et al. 2018). In photosynthetic tissues, photorespiration is the predominant serine biosynthesis route with 23% to 41% of photorespiratory serine leaving the cycle (Fu et al. 2023) to serve metabolic processes including sulfur-assimilation and

cysteine biosynthesis (Abadie and Tcherkez 2019). In this context, photorespiratory serine could produce *o*-acetyl serine required for sulfate-associated cysteine biosynthesis and subsequent ABA production.

So is the photorespiratory cycle a good target to improve stomatal opening/closure traits in the field? Chloroplast photorespiratory by-pass plants with improved photosynthesis might be expected to impact stomatal opening. That said, no significant changes were found in Arabidopsis (Maier et al. 2012) or tobacco (South et al. 2019), and only rice showed a higher *g<sub>s</sub>* (Nayak et al. 2022). Not all photorespiratory mutants exhibit altered *g<sub>s</sub>* or transpiration when transferred from high CO<sub>2</sub> to air (Dellero et al. 2015, 2016; Eisenhut et al. 2017) although antisense rice GOX showed a correlation between GOX activity and *g<sub>s</sub>* (Lu et al. 2014). Finally, reduction of end-of-day starch in *hpr1* leaves (Timm et al. 2012) was not associated with altered transpiration in the light (Eisenhut et al. 2017).


Many questions still remain unanswered such as how, where, and when does photorespiration impact stomatal opening/closing? It will be challenging but essential to answer these questions in the context of manipulating photorespiration to improve photosynthesis and plant yield since unwanted stomatal phenotypes could negatively impact water-use efficiency. Future directions to answer these questions should include determining stomatal kinetic responses to environmental cues in controlled atmospheric conditions differing in CO<sub>2</sub>/O<sub>2</sub> levels to modulate photorespiration in attached leaves. Other important avenues of investigation include determining whether photorespiration in guard cells, mesophyll cells, or both cell types is important using cell-specific expression lines, and identifying how photorespiration impacts metabolites involved in stomatal movements using single-cell metabolomics associated with stable-isotope labeling.

## What adaptive changes make an enzyme suitable for the C<sub>4</sub> carbon concentrating pathway, and how do critical and lineage-specific amino acid substitutions affect plant physiology?

(Written by Clarisa E. Alvarez, Veronica G. Maurino, and Marcos A. Tronconi)

### C<sub>4</sub> proteins have accumulated different types of adaptive changes

Photorespiration evolved as a selective response to Rubisco promiscuity, as a high-energy pathway to remove 2-phosphoglycolate, an unwanted reaction product. In C<sub>3</sub> plants, photorespiration negatively affects fitness under conditions that reduce the relative concentration of CO<sub>2</sub>, such as high temperature, drought, and salinity. Some land plants have evolved a biochemical CO<sub>2</sub> pump known as the C<sub>4</sub> carbon concentrating pathway or C<sub>4</sub> pathway. In the C<sub>4</sub> pathway, CO<sub>2</sub> is prefixed to form a C<sub>4</sub> acid, which is then decarboxylated in the vicinity of Rubisco. These reactions are usually spatially separated between two cell types (Fig. 11A) (Maier et al. 2011; Drincovich et al. 2016). By increasing the CO<sub>2</sub> concentration around Rubisco, C<sub>4</sub> plants gain a beneficial reduction in photorespiratory flux. The evolution of the C<sub>4</sub> pathway resulted in the coordinated acquisition of new anatomical and biochemical features (Sage and Zhu 2011; Alvarez and Maurino 2023). It is widely accepted that anatomical C<sub>4</sub> traits were acquired prior to C<sub>4</sub> biochemistry (Christin and Osborne 2014). On this basis, C<sub>4</sub> pathway genes were formed by co-opting copies of



**Figure 11.** Aspects of the  $C_4$  pathway. On the right: a schematic representation of the  $C_4$  biochemical pathway as found in maize and *Flaveria*. The main fluxes are indicated by thicker arrows. CBC, Calvin–Benson–Bassham Cycle. In the middle: 3D structural representations of tetrameric  $C_4$ -PEPC (bottom) and  $C_4$ -NADP-ME (top). On the left: subunits of  $C_4$ -PEPC and  $C_4$ -NADP-ME, highlighting differentially substituted amino acids. Amino acids in orange represent critical substitutions. Amino acids in black represent lineage-specific substitutions. The protein structures for *Sorghum bicolor*  $C_4$ -NADP-ME (PDB code 6C7N) and *Flaveria trinervia*  $C_4$ -PEPC (PDB code 3ZGE) were modeled using PyMOL Molecular Graphics System, version 2.0 Schrödinger, LLC.

duplicated genes that already existed in  $C_3$  species (Christin et al. 2013). In most cases, the same gene lineage has been selected repeatedly by independent  $C_4$  origins. The bias towards co-opted genes may be related to fewer key genetic changes in their regulatory and coding regions, which need to accumulate rapidly to acquire the  $C_4$  function (Copley 2020).

Some of the adaptive amino acid changes are critical for  $C_4$  function: if one of these is mutated back to the amino acid present in the  $C_3$  version of the enzyme, there is a loss of crucial  $C_4$  biochemical properties (kinetic, regulatory, and/or structural). In addition, other adaptive changes have occurred to optimize the  $C_4$  function, either (i) by counteracting perturbations in protein properties caused by previous critical  $C_4$  changes, such as a reduction in stability of the protein structure, or (ii) by adapting the enzyme activity to the new physiological environment that accompanies  $C_4$  activities. Candidates for critical adaptive amino acid changes can be identified through an analysis of protein sequences from different  $C_4$  lineages in the form of strictly differentially substituted positions (Alvarez et al. 2019; Hüdig et al. 2022). A strictly differentially substituted position is one where the protein sequences in the  $C_4$  species contain an identical amino acid, while a different amino acid is shared by all  $C_3$  sequences. The data obtained so far indicate that critical adaptive changes for  $C_4$  innovations have evolved with a high degree of convergence between distant  $C_4$  lineages, as in the case of PEPC (Christin et al. 2007; Rosnow et al. 2014; Lyu et al. 2021), or restricted to genetically close  $C_4$  lineages, as in the case of NADP-ME (Alvarez et al. 2019; Alvarez and Maurino 2023) and NAD-ME (Tronconi et al. 2020; Hüdig et al. 2022), demonstrating that alternative solutions to critical  $C_4$  requirements exist. Such sequence analyses also reveal a large number of molecular adaptations that are lineage-specific (Fig. 11, B and C).

### Innovation is easy, optimization is complicated: how can we analyze the impact of adaptive changes of $C_4$ proteins?

$C_4$  innovations can be achieved by a rather small number of critical changes, some of which evolved divergently while others

appear to have occurred convergently across  $C_4$  lineages. In this regard, when comparing the differentially substituted residues between the  $C_4$ -PEPC from Poaceae, Chenopodiaceae, and Asteraceae, which are genetically very distant, only one residue underwent an identical substitution in all three lineages. This substitution causes the  $C_4$ -PEPC to acquire affinity values for substrates similar to those of non- $C_4$  isoforms. This suggests that only minimal changes to the  $C_3$  ortholog may be required to perform the specialized  $C_4$  function. However, the presence of a large number of lineage-specific amino acid substitutions in  $C_4$  proteins indicates that the optimization of  $C_4$  function is much more complicated and may depend on lineage-specific physiological details. During the evolution of  $C_4$  biochemistry, the specific compartmentalization of enzyme activities and increased protein abundances created a new cellular environment (Lyu et al. 2021). In addition, the evolution of different biochemical and anatomical  $C_4$  subtypes imposed different physiological constraints between the  $C_4$  lineages. As a result,  $C_4$  enzymes coordinately adapted to the new metabolic context and the need for high carbon flux by fine-tuning their function through lineage-specific amino acid substitutions. An illustration of this process was provided by Lyu et al. (2021), who found that several genes encoding proteins associated with the  $C_4$  pathway in *Flaveria* species showed highly coordinated patterns of gene expression and modification in protein sequence.

The details of the molecular evolution of most  $C_4$  enzymes are still largely unknown. Present knowledge is based on the analysis of  $C_4$  enzymes from a small number of species in a few  $C_4$  lineages. Thus, it is not yet clear whether critical  $C_4$  changes are generally convergent, or whether they depend on the co-opted gene or on specific features of  $C_4$  subtypes. So, what adaptive amino acid substitutions make a protein suitable for  $C_4$  biochemistry in each lineage? To answer this question, much more data from more species is needed to allow intra- and inter-lineage analyses to associate critical  $C_4$  changes with specific amino acid replacements.

Data on the kinetic and regulatory properties of an enzyme are rarely collected under conditions that mimic physiological ones, and they are therefore not sufficient to fully understand an

enzyme's contribution to fitness. In fact, the performance of an enzyme depends on the physiological environment, emphasizing the importance of studying how adaptive changes affect plant physiology. At present, our knowledge of the in planta effects of adaptive changes is at a nascent stage. For example, in maize, *Amaranthus*, and *Flaveria*, the high activity level and low malate sensitivity of C<sub>4</sub>-PEPC were attributed to the phosphorylation state of the enzyme (Jiao and Chollet 1988; Tsuchida et al. 2001; Avasthi et al. 2011). However, the lack of phosphorylation of C<sub>4</sub>-PEPC from *Flaveria bidentis* was not crucial for PEPC activity or high photosynthesis rates under standard greenhouse conditions (Furumoto et al. 2007). Another example is the pH-dependent inhibition by malate of C<sub>4</sub>-NADP-ME from maize, sorghum (*Sorghum bicolor*), and *Setaria italica* (Alvarez et al. 2019). The question is, how does this regulatory property—exclusive to the C<sub>4</sub> isoform—affect plant physiology in a day-night cycle?

Several genetic engineering strategies to introduce a “C<sub>4</sub>-like pathway” into C<sub>3</sub> plants have failed to improve photosynthesis, most likely due to the lack of adaptation of the C<sub>4</sub> enzyme to the intracellular C<sub>3</sub> environment. There is a need to assess the biochemical and molecular characteristics of C<sub>4</sub> enzymes in planta to understand how and under what conditions specific adaptive changes may affect plant physiology. In the future, new technologies such as CRISPR will facilitate the introduction of wild-type and mutated versions of C<sub>4</sub> enzymes in C<sub>3</sub> and C<sub>4</sub> plants and the further analysis of their impact on plant metabolism and physiology. For example, to assess whether lineage-specific adaptive changes are related to the optimization of C<sub>4</sub> function, C<sub>4</sub> mutant plants lacking their own photosynthetic C<sub>4</sub>-enzyme but expressing the C<sub>4</sub> isoform of a distant lineage could be generated (e.g. C<sub>4</sub>-PEPC from grasses expressed in the C<sub>4</sub> *Flaveria trinervia* C<sub>4</sub>-*pepc*<sup>-</sup> mutant). In addition, C<sub>4</sub> mutant plants expressing a distant C<sub>4</sub> enzyme that contains amino acid substitutions specific to the lineage in which it is expressed could also be tested. These analyses would contribute to our overall understanding of the regulatory processes that are involved in the intricate dynamics of C<sub>4</sub> plant metabolism.

## Conclusions

The questions covered here span a wide range of processes related to oxygenic photosynthesis. A few common themes and conclusions can nonetheless be drawn from these diverse topics. First, protein interactions are critical and govern almost all cellular processes. The most exciting and important answers to unsolved questions will come from identifying the key proteins and their interacting partners, where and when the interactions take place, how they interact, and how they are regulated. A key aspect of many of the topics covered here is an emphasis on the importance of timing: when are molecular mechanisms active or interacting together? This is likely true for any biological system; therefore, answers to these questions will provide fundamental insights not only into photosynthesis but many other biochemical processes as well. Second, the core photosystems PSI and PSII are highly conserved among photosynthetic organisms, but many aspects of their regulation, from light harvesting to how the resulting chemical energy is dispersed and used in the cell and how excess energy is dissipated, are highly diversified across the plant kingdom. Therefore, more work is needed not only on the historical model systems (e.g. *Chlamydomonas*, *Arabidopsis*) but on developing new model systems and understanding the diversity of photosynthesis in all its forms. Our discussions of the above questions further suggest that we are entering an era of understanding—and being able to

interrogate—the ecophysiology of photosynthesis at the molecular level, which will be critical to understanding the diversity of photosynthetic regulatory mechanisms.

## Acknowledgments

The authors apologize to those whose contributions could not be cited due to editorial restrictions. The authors thank Patrice Salomé for his excellent suggestions and work to improve figures, especially Figs. 5, 6, 7, 9, and 10. J.Y., V.K.Y., and J.K. would like to thank all the students, postdoctoral fellows, and collaborators for their contributions to their research. A.R.G. thanks Adrien Burlacot for useful recommendations on the alternative electron flow section. Y.A., H.M., and L.N. thank Dr. Dmitry Shevela (ShevelaDesign AB, Sweden) for the graphical work. M.H. thanks past and present Metaboactions team members for their hard work and stimulating discussions throughout the years. C.E.A., V.G.M., and M.A.T. thank Daria Chrobok (DC SciArt, Sweden) for professional graphic work for Fig. 11.

## Author contributions

All authors contributed to writing and revising the article.

## Funding

Work in the lab of J.Y., V.K.Y., and J.K. was supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division (CSGB) (J.Y., V.K.Y., J.K.) for X-ray spectroscopy and crystallography data collection and analysis and methods development for photosynthetic systems, by the National Institutes of Health) Grants 1R35GM149528 (V.K.Y.) for PS II biochemical studies, GM110501 (J.Y.) and GM126289 (J.K.) for instrumentation development for XFEL experiments. Work by G.S.S.-C. and D.H. was supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division under Award #DE-SC0018097 to G.S.S.-C. Work by N.H. and C.B. was funded by the Deutsche Forschungsgemeinschaft (DFG, Bu812/8, Bu812/10) and supported by Horizon 2020 of the European Union Research and Innovation Program ITN (grant no. 675006). The financial support of a UK Research and Innovation Future Leaders Fellowship (MR/T017546/1 and MR/Y011635/1 to T.D.K. and T.C.) and an EPSRC Doctoral Training Partnership Scholarship (to E.C.) is gratefully acknowledged. Relevant work in the C.W.M. lab is supported by BBSRC (BB/W001012/1). Relevant work by W.W. and B.D.E. was supported by a Human Frontier Science Program Grant (RGP0005/2021) and ERC consolidator grant “cryOcean” (fulfilled by the Swiss State Secretariat for Education, Research and Innovation, M822.00045). Much of the work performed in the lab of A.R.G. on the export of metabolites from the chloroplast and alternative electron transport pathways was supported by U.S. Department of Energy award DE-SC0019417.

Y.A. acknowledges Academy of Finland (project no. 315119). A.B. acknowledges the support of the Carnegie Institution for Science. Work in the team of M.H. has been supported by the Agence Nationale de la Recherche (ANR) (ANR-10-LABX-0040-SPS, ANR-11-IDEX-0003-02, ANR-14-CE19-0015 REGUL3P), the CNRS, INRAE, University Paris-Saclay and the Saclay Plant Sciences Graduate School of Research. V.G.M. was funded by the Deutsche Forschungsgemeinschaft (DFG grant

MA2379/20-1 project number 441941117), and C.E.A. was funded by the ANPCyT grant PICT-2019-00079.

Conflict of interest statement. None declared.

## References

- Abadie C, Tcherkez G. Plant sulfur metabolism is stimulated by photorespiration. *Comm Biol*. 2019;2(1):379. <https://doi.org/10.1038/s42003-019-0616-y>
- Alboresi A, Storti M, Cendron L, Morosinotto T. Role and regulation of class-C flavodiiron proteins in photosynthetic organisms. *Biochem J*. 2019a;476(17):2487–2498. <https://doi.org/10.1042/BCJ20180648>
- Alboresi A, Storti M, Morosinotto T. Balancing protection and efficiency in the regulation of photosynthetic electron transport across plant evolution. *New Phytol*. 2019b;221(1):105–109. <https://doi.org/10.1111/nph.15372>
- Allahverdiyeva Y, Isojärvi J, Zhang P, Aro EM. Cyanobacterial oxygenic photosynthesis is protected by flavodiiron proteins. *Life (Basel)*. 2015a;5(1):716–743. <https://doi.org/10.3390/life5010716>
- Allahverdiyeva Y, Mustila H, Ermakova M, Bersanini L, Richaud P, Ajlani G, Battchikova N, Cournac L, Aro EM. Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. *Proc Natl Acad Sci U S A*. 2013;110(10):4111–4116. <https://doi.org/10.1073/pnas.1221194110>
- Allahverdiyeva Y, Suorsa M, Tikkanen M, Aro EM. Photoprotection of photosystems in fluctuating light intensities. *J Exp Bot*. 2015b;66(9):2427–2436. <https://doi.org/10.1093/jxb/eru463>
- Allen JF, Forsberg J. Molecular recognition in thylakoid structure and function. *Trends Plant Sci*. 2001;6(7):317–326. [https://doi.org/10.1016/S1360-1385\(01\)02010-6](https://doi.org/10.1016/S1360-1385(01)02010-6)
- Allorant G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D, Petroustos D, Kuntz M, Breyton C, et al. A dual strategy to cope with high light in *Chlamydomonas reinhardtii*. *Plant Cell*. 2013;25(2):545–557. <https://doi.org/10.1105/tpc.112.108274>
- Alvarez CE, Bovdilova A, Hoppner A, Wolff CC, Saigo M, Trajtenberg F, Zhang T, Buschiazzo A, Nagel-Steger L, Drincovich MF, et al. Molecular adaptations of NADP-malic enzyme for its function in C4 photosynthesis in grasses. *Nat Plants*. 2019;5(7):755–765. <https://doi.org/10.1038/s41477-019-0451-7>
- Alvarez CE, Maurino VG. Adaptive diversity in structure and function of C4 photosynthetic components. *Biochem Soc Trans*. 2023;51(3):1157–1168. <https://doi.org/10.1042/BST20221279>
- Andersson B, Anderson JM. Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *Biochim Biophys Acta*. 1980;593(2):427–440. [https://doi.org/10.1016/0005-2728\(80\)90078-X](https://doi.org/10.1016/0005-2728(80)90078-X)
- Appel J, Hueren V, Boehm M, Gutekunst K. Cyanobacterial in vivo solar hydrogen production using a photosystem I-hydrogenase (PsaD-HoxYH) fusion complex. *Nature Energy*. 2020;5(6):458–467. <https://doi.org/10.1038/s41560-020-0609-6>
- Armbruster U, Leonelli L, Correa Galvis V, Strand D, Quinn EH, Jonikas MC, Niyogi KK. Regulation and levels of the thylakoid K<sup>+</sup>/H<sup>+</sup> antiporter KEA3 shape the dynamic response of photosynthesis in fluctuating light. *Plant Cell Physiol*. 2016;57(7):1557–1567. <https://doi.org/10.1093/pcp/pcw085>
- Arshad R, Calvaruso C, Boekema E, Büchel C, Kouril R. Revealing the architecture of the photosynthetic apparatus in the diatom *Thalassiosira pseudonana*. *Plant Physiol*. 2021;186(4):2124–2136. <https://doi.org/10.1093/plphys/kiab208>
- Asada K. THE WATER-WATER CYCLE IN CHLOROPLASTS: scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol Plant Mol Biol*. 1999;50(1):601–639. <https://doi.org/10.1146/annurev.arplant.50.1.601>
- Avasthi UK, Izui K, Raghavendra AS. Interplay of light and temperature during the in planta modulation of C4 phosphoenolpyruvate carboxylase from the leaves of *Amaranthus hypochondriacus* L.: diurnal and seasonal effects manifested at molecular levels. *J Exp Bot*. 2011;62(3):1017–1026. <https://doi.org/10.1093/jxb/erq333>
- Bailleul B, Rogato A, De Martino A, Coesel S, Cardol P, Bowler C, Falciatore A, Finazzi G. An atypical member of the light-harvesting complex stress-related protein family modulates diatom responses to light. *Proc Natl Acad Sci U S A*. 2010;107(42):18214–18219. <https://doi.org/10.1073/pnas.1007703107>
- Barber J, Andersson B. Too much of a good thing: light can be bad for photosynthesis. *Trends Biochem Sci*. 1992;17(2):61–66. [https://doi.org/10.1016/0968-0004\(92\)90503-2](https://doi.org/10.1016/0968-0004(92)90503-2)
- Barros T, Kühlbrandt W. Crystallisation, structure and function of plant light-harvesting Complex II. *Biochim Biophys Acta*. 2009;1787(6):753–772. <https://doi.org/10.1016/j.bbabi.2009.03.012>
- Batool S, Uslu VV, Rajab H, Ahmad N, Waadt R, Geiger D, Malagoli M, Xiang CB, Hedrich R, Rennenberg H, et al. Sulfate is incorporated into cysteine to trigger ABA production and stomata closure. *Plant Cell*. 2018;30(12):2973–2987. <https://doi.org/10.1105/tpc.18.00612>
- Bersanini L, Allahverdiyeva Y, Battchikova N, Heinz S, Lespinasse M, Ruohisto E, Mustila H, Nickelsen J, Vass I, Aro EM. Dissecting the photoprotective mechanism encoded by the *flv4-2* operon: a distinct contribution of Sll0218 in photosystem II stabilization. *Plant Cell Environ*. 2017;40(3):378–389. <https://doi.org/10.1111/pce.12872>
- Bersanini L, Battchikova N, Jokel M, Rehman A, Vass I, Allahverdiyeva Y, Aro EM. Flavodiiron protein Flv2/Flv4-related photoprotective mechanism dissipates excitation pressure of PSII in cooperation with phycobilisomes in Cyanobacteria. *Plant Physiol*. 2014;164(2):805–818. <https://doi.org/10.1104/pp.113.231969>
- Bhattacharya D, Medlin L. The phylogeny of plastids: a review based on comparisons of small-subunit ribosomal RNA coding regions. *J Phycol*. 1995;31(4):489–498. <https://doi.org/10.1111/j.1529-8817.1995.tb02542.x>
- Bhowmick A, Hussein R, Bogacz I, Simon PS, Ibrahim M, Chatterjee R, Doyle MD, Cheah MH, Fransson T, Chernev P, et al. Structural evidence for intermediates during O<sub>2</sub> formation in photosystem II. *Nature*. 2023;617(7961):629–636. <https://doi.org/10.1038/s41586-023-06038-z>
- Bockwoldt M, Heiland I, Fischer K. The evolution of the plastid phosphate translocator family. *Planta*. 2019;250(1):245–261. <https://doi.org/10.1007/s00425-019-03161-y>
- Brown KA, Guo Z, Tokmina-Lukaszewska M, Scott LW, Lubner CE, Smolinski S, Mulder DW, Bothner B, King PW. The oxygen reduction reaction catalyzed by *Synechocystis* sp. PCC 6803 flavodiiron proteins. *Sustain Energ Fuels*. 2019;3(11):3191–3200. <https://doi.org/10.1039/c9se00523d>
- Büchel C. Evolution and function of light harvesting proteins. *J Plant Physiol*. 2015;172(19):62–75. <https://doi.org/10.1016/j.jplph.2014.04.018>
- Buchert F, Hamon M, Gabelein P, Scholz M, Hippler M, Wollman FA. The labile interactions of cyclic electron flow effector proteins. *J Biol Chem*. 2018;93(45):17559–17573. <https://doi.org/10.1074/jbc.RA118.004475>
- Buck JM, Kroth PG, Lepetit B. Identification of sequence motifs in LhcX proteins that confer qE-based photoprotection in the diatom *Phaeodactylum tricornutum*. *Plant J*. 2021;108(6):1721–1734. <https://doi.org/10.1111/tpj.15539>
- Burlacot A, Richaud P, Gosset A, Li-Beisson Y, Peltier G. Algal photosynthesis converts nitric oxide into nitrous oxide. *Proc Natl Acad Sci U S A*. 2019;116(12):5845–5850. <https://doi.org/10.1073/pnas.1812111116>

- Sci U S A. 2020;117(5):2704–2709. <https://doi.org/10.1073/pnas.1915276117>
- Butkus V, Gelzinis A, Augulis R, Gall A, Büchel C, Robert B, Zigmantas D, Valkunas L, Abramavicius D. Coherence and population dynamics of chlorophyll excitations in FCP complex: two-dimensional spectroscopy study. *J Chem Phys.* 2015;142(21):212414. <https://doi.org/10.1063/1.4914098>
- Calvaruso C, Rokka A, Aro E-M, Büchel C. Specific Lhc proteins are bound to PSI or PSII supercomplexes in the diatom *Thalassiosira pseudonana*. *Plant Physiol.* 2020;183(1):67–79. <https://doi.org/10.1104/pp.20.00042>
- Cantrell M, Peers G. A mutant of *Chlamydomonas* without LHCSR maintains high rates of photosynthesis, but has reduced cell division rates in sinusoidal light conditions. *PLoS One.* 2017;12(6):e0179395. <https://doi.org/10.1371/journal.pone.0179395>
- Cardol P, Alric J, Girard-Bascou J, Franck F, Wollman FA, Finazzi G. Impaired respiration discloses the physiological significance of state transitions in *Chlamydomonas*. *Proc Natl Acad Sci U S A.* 2009;106(37):15979–15984. <https://doi.org/10.1073/pnas.0908111106>
- Carmo-Silva E, Sharwood RE. Rubisco and its regulation—major advances to improve carbon assimilation and productivity. *J Exp Bot.* 2023;74(2):507–509. <https://doi.org/10.1093/jxb/erac475>
- Casella S, Huang F, Mason D, Zhao G-Y, Johnson GN, Mullineaux CW, Liu L-N. Dissecting the native architecture and dynamics of cyanobacterial photosynthetic machinery. *Mol Plant.* 2017;10(11):1434–1448. <https://doi.org/10.1016/j.molp.2017.09.019>
- Charuvi D, Nevo R, Kaplan-Ashiri I, Shimoni E, Reich Z. Studying the supramolecular organization of photosynthetic membranes within freeze-fractured leaf tissues by cryo-scanning electron microscopy. *J Vis Exp.* 2016(112):e54066. <https://doi.org/10.3791/54066>
- Charuvi D, Nevo R, Shimoni E, Naveh L, Zia A, Adam Z, Farrant JM, Kirchoff H, Reich Z. Photoprotection conferred by changes in photosynthetic protein levels and organization during dehydration of a homoiochlorophyllous resurrection plant. *Plant Physiol.* 2015;167(4):1554–1565. <https://doi.org/10.1104/pp.114.255794>
- Chaux F, Burlacot A, Mekhalfi M, Auroy P, Blangy S, Richaud P, Peltier G. Flavodiiron proteins promote fast and transient O<sub>2</sub> photoreduction in *Chlamydomonas*. *Plant Physiol.* 2017;174(3):1825–1836. <https://doi.org/10.1104/pp.17.00421>
- Chernev P, Fischer S, Hoffmann J, Oliver N, Assunção R, Yu B, Burnap RL, Zaharieva I, Nürnberg DJ, Haumann M, et al. Light-driven formation of manganese oxide by today's photosystem II supports evolutionarily ancient manganese-oxidizing photosynthesis. *Nat Commun.* 2020;11(1):6110. <https://doi.org/10.1038/s41467-020-19852-0>
- Cho C, Ishii R, Hyeon S-B, Suzuki A. Stomatal regulation by aminoacetoneitrile, a photorespiration inhibitor. *Plant Cell Physiol.* 1987;28:1407–1410. <https://doi.org/10.1093/oxfordjournals.pcp.a077432>
- Christin PA, Boxall SF, Gregory R, Edwards EJ, Hartwell J, Osborne CP. Parallel recruitment of multiple genes into c4 photosynthesis. *Genome Biol. Evol.* 2013;5(11):2174–2187. <https://doi.org/10.1093/gbe/evt168>
- Christin PA, Osborne CP. The evolutionary ecology of C4 plants. *New Phytol.* 2014;204(4):765–781. <https://doi.org/10.1111/nph.13033>
- Christin PA, Salamin N, Savolainen V, Duvall MR, Besnard G. C4 photosynthesis evolved in grasses via parallel adaptive genetic changes. *Curr Biol.* 2007;17(14):1241–1247. <https://doi.org/10.1016/j.cub.2007.06.036>
- Clausen CH, Brooks MD, Li T-D, Grob P, Kemalyan G, Nogales E, Niyogi KK, Fletcher DA. Dynamic mechanical responses of *Arabidopsis* thylakoid membranes during PSII-specific illumination. *Biophys J.* 2014;106(9):1864–1870. <https://doi.org/10.1016/j.bpj.2014.03.016>
- Copley SD. Evolution of new enzymes by gene duplication and divergence. *FEBS J.* 2020;287(7):1262–1283. <https://doi.org/10.1111/febs.15299>
- Croce R, Carmo-Silva E, Cho YB, Ermakova M, Harbinson J, Lawson T, McCormick AJ, Niyogi KK, Ort DR, Patel-Tupper D. Perspectives on improving photosynthesis to increase crop yield. *Plant Cell.* 2024;36(10):3944–3973. <https://doi.org/10.1093/plcell/koae132>
- Croce R, van Amerongen H. Light harvesting in oxygenic photosynthesis: structural biology meets spectroscopy. *Science.* 2020;369(6506):eaay2058. <https://doi.org/10.1126/science.aay2058>
- Cruz JA, Savage LJ, Zegarac R, Hall CC, Satoh-Cruz M, Davis GA, Kovac WK, Chen J, Kramer DM. Dynamic environmental photosynthetic imaging reveals emergent phenotypes. *Cell Syst.* 2016;2(6):365–377. <https://doi.org/10.1016/j.cels.2016.06.001>
- Dai L, Tan LM, Jiang YL, Shi Y, Wang P, Zhang JP, Otomo ZY. Orientation assignment of LH2 and LH1-RC complexes from Thermochromatium tepidum reconstituted in PC liposome and their ultrafast excitation dynamics comparison between in artificial and in natural chromatophores. *Chem Phys Lett.* 2018;705:78–84. <https://doi.org/10.1016/j.cplett.2018.05.043>
- Dalcorso G, Pesaresi P, Masiero S, Aseeva E, Schunemann D, Finazzi G, Joliet P, Barbato R, Leister D. A complex containing PGR1 and PGR5 is involved in the switch between linear and cyclic electron flow in *Arabidopsis*. *Cell.* 2008;132(2):273–285. <https://doi.org/10.1016/j.cell.2007.12.028>
- Dang KV, Plet J, Tolleter D, Jokel M, Cuine S, Carrier P, Auroy P, Richaud P, Johnson X, Alric J, et al. Combined increases in mitochondrial cooperation and oxygen photoreduction compensate for deficiency in cyclic electron flow in *Chlamydomonas reinhardtii*. *Plant Cell.* 2014;26(7):3036–3050. <https://doi.org/10.1105/tpc.114.126375>
- Dao O, Kuhnert F, Weber APM, Peltier G, Li-Beisson Y. Physiological functions of malate shuttles in plants and algae. *Trends Plant Sci.* 2022;27(5):488–501. <https://doi.org/10.1016/j.tplants.2021.11.007>
- Dau H, Haumann M. The manganese complex of photosystem II in its reaction cycle—basic framework and possible realization at the atomic level. *Coord Chem Rev.* 2008;252(3–4):273–295. <https://doi.org/10.1016/j.ccr.2007.09.001>
- Daum B, Nicastro D, Austin J 2nd, McIntosh JR, Kühlbrandt W. Arrangement of photosystem II and ATP synthase in chloroplast membranes of spinach and pea. *Plant Cell.* 2010;22(4):1299–1312. <https://doi.org/10.1105/tpc.109.071431>
- Debus RJ. FTIR studies of metal ligands, networks of hydrogen bonds, and water molecules near the active site Mn<sub>4</sub>CaO<sub>5</sub> cluster in photosystem II. *Biochim Biophys Acta.* 2015;1847(1):19–34. <https://doi.org/10.1016/j.bbabi.2014.07.007>
- Dekker JP, Boekema EJ. Supramolecular organization of thylakoid membrane proteins in green plants. *Biochim Biophys Acta.* 2005;1706(1–2):12–39. <https://doi.org/10.1016/j.bbabi.2004.09.009>
- Dellero Y, Jossier M, Glab N, Oury C, Tcherkez G, Hodges M. Decreased glycolate oxidase activity leads to altered carbon allocation and leaf senescence after a transfer from high CO<sub>2</sub> to ambient air in *Arabidopsis thaliana*. *J Exp Bot.* 2016;67(10):3149–3163. <https://doi.org/10.1093/jxb/erw054>
- Dellero Y, Lamothe-Sibold M, Jossier M, Hodges M. *Arabidopsis thaliana ggt1* photorespiratory mutants maintain leaf carbon/nitrogen balance by reducing RuBisCO content and plant growth. *Plant J.* 2015;83(6):1005–1118. <https://doi.org/10.1111/tpj.12945>
- Depège N, Bellafiore S, Rochaix J-D. Role of chloroplast protein kinase stt7 in LHCI phosphorylation and state transition in

- Chlamydomonas. *Science*. 2003;299(5612):1572–1575. <https://doi.org/10.1126/science.1081397>
- De Souza AP, Burgess SJ, Doran L, Hansen J, Manukyan L, Maryn N, Gotarkar D, Leonelli L, Niyogi KK, Long SP. Soybean photosynthesis and crop yield are improved by accelerating recovery from photoprotection. *Science*. 2022;377(6608):851–854. <https://doi.org/10.1126/science.adc9831>
- Desplats C, Mus F, Cuine S, Billon E, Cournac L, Peltier G. Characterization of *nda2*, a plastoquinone-reducing type II NAD(P)H dehydrogenase in *Chlamydomonas* chloroplasts. *J Biol Chem*. 2009;284(7):4148–4157. <https://doi.org/10.1074/jbc.M804546200>
- Dietzel L, Bräutigam K, Pfannschmidt T. Photosynthetic acclimation: state transitions and adjustment of photosystem stoichiometry—functional relationships between short-term and long-term light quality acclimation in plants. *FEBS J*. 2008;275(6):1080–1088. <https://doi.org/10.1111/j.1742-4658.2008.06264.x>
- Drincovich MF, Voll LM, Maurino VG. Editorial: on the diversity of roles of organic acids. *Front Plant Sci*. 2016;7:1592. <https://doi.org/10.3389/fpls.2016.01592>
- Duminil P. 2019. Characterization of two primary metabolism enzymes in *Arabidopsis thaliana*: phosphoglycerate mutase and phosphoglycolate phosphatase [PhD Dissertation] University Paris-Saclay.
- Durnford DG, Price JA, McKim SM, Sarchfield ML. Light-harvesting complex gene expression is controlled by both transcriptional and post-transcriptional mechanisms during photoacclimation in *Chlamydomonas reinhardtii*. *Phys Plantarum*. 2003;118(2):193–205. <https://doi.org/10.1034/j.1399-3054.2003.00078.x>
- Eberhard S, Finazzi G, Wollman FA. The dynamics of photosynthesis. *Ann Rev Genet*. 2008;42(1):463–515. <https://doi.org/10.1146/annurev.genet.42.110807.091452>
- Eicks M, Maurino V, Knappe S, Flugge UI, Fischer K. The plastidic pentose phosphate translocator represents a link between the cytosolic and the plastidic pentose phosphate pathways in plants. *Plant Physiol*. 2002;128(2):512–522. <https://doi.org/10.1104/pp.010576>
- Eisenhut M, Bräutigam A, Timm S, Florian A, Tohge T, Fernie AR, Bauwe H, Weber APM. Photorespiration is crucial for dynamic response of photosynthetic metabolism and stomatal movement to altered CO<sub>2</sub> availability. *Mol Plant*. 2017;10(1):47–61. <https://doi.org/10.1016/j.molp.2016.09.011>
- Eisenhut M, Georg J, Klähn S, Sakurai I, Mustila H, Zhang P, Hess WR, Aro EM. The antisense RNA *as1\_flv4* in the cyanobacterium *Synechocystis* sp. PCC 6803 prevents premature expression of the *flv4-2* operon upon shift in inorganic carbon supply. *J Biol Chem*. 2012;287(40):33153–33162. <https://doi.org/10.1074/jbc.M112.391755>
- Eisenhut M, Roell M, Weber APM. Mechanistic understanding of photorespiration paves the way to a new green revolution. *New Phytol*. 2019;223(4):1762–1769. <https://doi.org/10.1111/nph.15872>
- Engel BD, Schaffer M, Kuhn Cuellar L, Villa E, Plitzko JM, Baumeister W. Native architecture of the *Chlamydomonas* chloroplast revealed by in situ cryo-electron tomography. *Elife*. 2015;4:e04889. <https://doi.org/10.7554/eLife.04889>
- Ermakova M, Battchikova N, Allahverdiyeva Y, Aro EM. Novel heterocyst-specific flavodiiron proteins in *Anabaena* sp. PCC 7120. *FEBS Lett*. 2013;587(1):82–87. <https://doi.org/10.1016/j.febslet.2012.11.006>
- Ermakova M, Battchikova N, Richaud P, Leino H, Kosourov S, Isojärvi J, Peltier G, Flores E, Cournac L, Allahverdiyeva Y, et al. Heterocyst-specific flavodiiron protein Flv3B enables oxic diazotrophic growth of the filamentous cyanobacterium *Anabaena* sp. PCC 7120. *Proc Natl Acad Sci USA*. 2014;111(30):11205–11210. <https://doi.org/10.1073/pnas.1407327111>
- Faries KM, Kressel LL, Wander MJ, Holten D, Laible PD, Kirmaier C, Hanson DK. High throughput engineering to revitalize a vestigial electron transfer pathway in bacterial photosynthetic reaction centers. *J Biol Chem*. 2012;287(11):8507–8514. <https://doi.org/10.1074/jbc.M111.326447>
- Feng L, Raza MA, Li Z, Chen Y, Khalid MHB, Du J, Liu W, Wu X, Song C, Yu L, et al. The influence of light intensity and leaf movement on photosynthesis characteristics and carbon balance of soybean. *Front Plant Sci*. 2019;9:1952. <https://doi.org/10.3389/fpls.2018.01952>
- Feng Y, Li Z, Li X, Shen L, Liu X, Zhou C, Zhang J, Sang M, Han G, Yang W, et al. Structure of a diatom photosystem II supercomplex containing a member of lhcx family and dimeric FCP II. *Sci Adv*. 2023;9(43):eadi8446. <https://doi.org/10.1126/sciadv.adi8446>
- Fiebig OC, Harris D, Wang D, Hoffmann M, Schlaue-Cohen GS. Ultrafast dynamics of photosynthetic light harvesting: strategies for acclimation across organisms. *Annu Rev Phys Chem*. 2023;74(1):493–520. <https://doi.org/10.1146/annurev-physchem-083122-111318>
- Field CB, Behrenfeld MJ, Randerson JT, Falkowski P. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science*. 1998;281(5374):237–240. <https://doi.org/10.1126/science.281.5374.237>
- Fischer K, Kammerer B, Gutensohn M, Arbinger B, Weber A, Hausler RE, Flugge UI. A new class of plastidic phosphate translocators: a putative link between primary and secondary metabolism by the phosphoenolpyruvate/phosphate antiporter. *Plant Cell*. 1997;199:453–462. <https://doi.org/10.1105/tpc.9.3.453>
- Fleming GR, Schlaue-Cohen GS, Amarnath K, Zaks J. Design principles of photosynthetic light-harvesting. *Faraday Discuss*. 2012;155:27–41. <https://doi.org/10.1039/C1FD00078K>
- Flügel F, Timm S, Arrivault S, Florian A, Stitt M, Fernie AR, Bauwe H. The photorespiratory metabolite 2-phosphoglycolate regulates photosynthesis and starch accumulation in *Arabidopsis*. *Plant Cell*. 2017;29(10):2537–2551. <https://doi.org/10.1105/tpc.17.00256>
- Flügge U-I, Häusler RE, Ludewig F, Fischer K. Functional genomics of phosphate antiport systems of plastids. *Physiol Plant*. 2003;118(4):475–482. <https://doi.org/10.1034/j.1399-3054.2003.00137.x>
- Foudree A, Putarjunan A, Kambakam S, Nolan T, Fussell J, Pogorelko G, Rodermel S. The mechanism of variegation in immutans provides insight into chloroplast biogenesis. *Front Plant Sci*. 2012;3:260. <https://doi.org/10.3389/fpls.2012.00260>
- Fu X, Gregory LM, Weise SE, Walker BJ. Integrated flux and pool size analysis in plant central metabolism reveals unique roles of glycine and serine during photorespiration. *Nat Plants*. 2023;9(1):169–178. <https://doi.org/10.1038/s41477-022-01294-9>
- Furumoto T, Izui K, Quinn V, Furbank RT, von Caemmerer S. Phosphorylation of phosphoenolpyruvate carboxylase is not essential for high photosynthetic rates in the C4 species *Flaveria bidentis*. *Plant Physiol*. 2007;144(4):1936–1945. <https://doi.org/10.1104/pp.107.102541>
- Gan F, Zhang S, Rockwell NC, Martin SS, Lagarias JC, Bryant DA. Extensive remodeling of a cyanobacterial photosynthetic apparatus in far-red light. *Science*. 2014;345(6202):1312–1317. <https://doi.org/10.1126/science.1256963>
- Gehlen J, Panstruga R, Smets H, Merkelbach S, Kleines M, Porsch P, Fladung M, Becker I, Rademacher T, Häusler RE, et al. Effects of altered phosphoenolpyruvate carboxylase activities on transgenic C3 plant *Solanum tuberosum*. *Plant Mol Biol*. 1996;32(5):831–848. <https://doi.org/10.1007/BF00020481>

- Gerotto C, Alboresi A, Meneghesso A, Jokel M, Suorsa M, Aro EM, Morosinotto T. Flavodiiron proteins act as safety valve for electrons in *Physcomitrella patens*. *Proc Natl Acad Sci U S A*. 2016;113(43):12322–12327. <https://doi.org/10.1073/pnas.1606685113>
- Ghazaryan A, Akhtar P, Garab G, Lambrev PH, Büchel C. Involvement of the LhcX protein Fcp6 of the diatom *Cyclotella meneghiniana* in the macro-organization and structural flexibility of thylakoid membranes. *BBA-Bioenergetics*. 2016;1857(9):1373–1379. <https://doi.org/10.1016/j.bbabi.2016.04.288>
- Gisriel CJ, Bryant DA, Brudvig GW, Cardona T. Molecular diversity and evolution of far-red light-acclimated photosystem I. *Front Plant Sci*. 2023;14:1289199. <https://doi.org/10.3389/fpls.2023.1289199>
- Glatzel P, Bergmann U, Yano J, Visser H, Robblee JH, Gu WW, de Groot FMF, Christou G, Pecoraro VL, Cramer SP, et al. The electronic structure of mn in oxides, coordination complexes, and the oxygen-evolving complex of photosystem II studied by resonant inelastic X-ray scattering. *J Am Chem Soc*. 2004;126(32):9946–9959. <https://doi.org/10.1021/ja038579z>
- Glatzel P, Schroeder H, Pushkar Y, Boron T III, Mukherjee S, Christou G, Pecoraro VL, Messinger J, Yachandra VK, Bergmann U, et al. Electronic structural changes of mn in the oxygen-evolving complex of photosystem II during the catalytic cycle. *Inorg Chem*. 2013;52(10):5642–5644. <https://doi.org/10.1021/ic4005938>
- Goral TK, Johnson MP, Brain APR, Kirchoff H, Ruban AV, Mullineaux CW. Visualizing the mobility and distribution of chlorophyll proteins in higher plant thylakoid membranes: effects of photoinhibition and protein phosphorylation. *Plant J*. 2010;62(6):948–959. <https://doi.org/10.1111/j.0960-7412.2010.04207.x>
- Goral TK, Johnson M, Duffy CDP, Brain APR, Ruban AV, Mullineaux CW. Light-harvesting antenna composition controls the macrostructure and dynamics of thylakoid membranes in *Arabidopsis*. *Plant J*. 2012;69(2):289–301. <https://doi.org/10.1111/j.1365-3113.2011.04790.x>
- Greife P, Schönborn M, Capone M, Assunção R, Narzi D, Guidoni L, Dau H. The electron–proton bottleneck of photosynthetic oxygen evolution. *Nature*. 2023;617(7961):623–628. <https://doi.org/10.1038/s41586-023-06008-5>
- Grouneva I, Jakob T, Wilhelm C, Goss R. A new multicomponent NPQ mechanism in the diatom *Cyclotella meneghiniana*. *Plant Cell Physiol*. 2008;49(8):1217–1225. <https://doi.org/10.1093/pcp/pcn097>
- Gundermann K, Büchel C. Factors determining the fluorescence yield of fucoxanthin-chlorophyll complexes (FCP) involved in non-photochemical quenching in diatoms. *BBA-Bioenergetics*. 2012;1817(7):1044–1052. <https://doi.org/10.1016/j.bbabi.2012.03.008>
- Gundermann K, Schmidt M, Weisheit W, Mittag M, Büchel C. Identification of several sub-populations in the pool of light harvesting proteins in the pennate diatom *Phaeodactylum tricorutum*. *BBA-Bioenergetics*. 2013;1827(3):303–310. <https://doi.org/10.1016/j.bbabi.2012.10.017>
- Guo J, Nguyen AY, Dai Z, Su D, Gaffrey MJ, Moore RJ, Jacobs JM, Monroe ME, Smith RD, Koppelaar DW, et al. Proteome-wide light/dark modulation of thiol oxidation in cyanobacteria revealed by quantitative site-specific redox proteomics. *Mol Cell Proteomics*. 2014;13(12):3270–3285. <https://doi.org/10.1074/mcp.M114.041160>
- Harris D, Toporik H, Schlau-Cohen GS, Mazor Y. Energetic robustness to large scale structural fluctuations in a photosynthetic supercomplex. *Nat Commun*. 2023;14(1):4650. <https://doi.org/10.1038/s41467-023-40146-8>
- Hausler RE, Schlieben NH, Schulz B, Flugge UI. Compensation of decreased triose phosphate/phosphate translocator activity by accelerated starch turnover and glucose transport in transgenic tobacco. *Planta*. 1998;204(3):366–376. <https://doi.org/10.1007/s004250050268>
- He C, Berkowitz O, Hu S, Zhao Y, Qian K, Shou H, Whelan J, Wang J. Co-regulation of mitochondrial and chloroplast function: molecular components and mechanisms. *Plant Comm*. 2023;4(1):100496. <https://doi.org/10.1016/j.xplc.2022.100496>
- He S, Chou HT, Matthies D, Wunder T, Meyer MT, Atkinson N, Martinez-Sanchez A, Jeffrey PD, Port SA, Patena W, et al. The structural basis of Rubisco phase separation in the pyrenoid. *Nat Plants*. 2020;6(12):1480–1490. <https://doi.org/10.1038/s41477-020-00811-y>
- Helman Y, Tchernov D, Reinhold L, Shibata M, Ogawa T, Schwarz R, Ohad I, Kaplan A. Genes encoding A-type flavoproteins are essential for photoreduction of O<sub>2</sub> in cyanobacteria. *Curr Biol*. 2003;13(3):230–235. [https://doi.org/10.1016/s0960-9822\(03\)00046-0](https://doi.org/10.1016/s0960-9822(03)00046-0)
- Herdean A, Teardo E, Nilsson AK, Pfeil BE, Johansson ON, Ünneper R, Nagy G, Zsiros O, Dana S, Solymosi K, et al. A voltage-dependent chloride channel fine-tunes photosynthesis in plants. *Nat Commun*. 2016;7(1):11654. <https://doi.org/10.1038/ncomms11654>
- Horrer D, Flüttsch S, Pazmino D, Matthews JS, Thalmann M, Nigro A, Leonhardt N, Lawson T, Santelia D. Blue light induces a distinct starch degradation pathway in guard cells for stomatal opening. *Curr Biol*. 2016;26(3):362–370. <https://doi.org/10.1016/j.cub.2015.12.036>
- Houille-Vernes L, Rappaport F, Wollman FA, Alric J, Johnson X. Plastid terminal oxidase 2 (PTOX2) is the major oxidase involved in chlororespiration in *Chlamydomonas*. *Proc Natl Acad Sci U S A*. 2011;108(51):20820–20825. <https://doi.org/10.1073/pnas.1110518109>
- Huang W, Krishnan A, Plett A, Meagher M, Linka N, Wang Y, Ren B, Findinier J, Redekop P, Fakhima N, et al. *Chlamydomonas* mutants lacking chloroplast TRIOSE PHOSPHATE TRANSPORTER3 are metabolically compromised and light sensitive. *Plant Cell*. 2023;35(7):2592–2614. <https://doi.org/10.1093/plcell/koad095>
- Hüdig M, Tronconi MA, Zubimendi JP, Sage TL, Poschmann G, Bickel D, Gohlke H, Maurino VG. Respiratory and C<sub>4</sub>-photosynthetic NAD-malic enzyme coexist in bundle sheath cell mitochondria and evolved via association of differentially adapted subunits. *Plant Cell*. 2022;34(1):597–615. <https://doi.org/10.1093/plcell/koab265>
- Huokko T, Ni T, Dykes GF, Simpson DM, Brownridge P, Conradi FD, Beynon RJ, Nixon PJ, Mullineaux CW, Zhang P, et al. Probing the biogenesis pathway and dynamics of thylakoid membranes. *Nat Commun*. 2021;12(1):3475. <https://doi.org/10.1038/s41467-021-23680-1>
- Hussein R, Ibrahim M, Bhowmick A, Simon PS, Bogacz I, Doyle MD, Dobbek H, Zouni A, Messinger J, Yachandra VK, et al. Evolutionary diversity of proton and water channels on the oxidizing side of photosystem II and their relevance to function. *Photosynth. Res*. 2023;158(2):91–107. <https://doi.org/10.1007/s11120-023-01018-w>
- Hussein R, Ibrahim M, Bhowmick A, Simon PS, Chatterjee R, Lassalle L, Doyle M, Bogacz I, Kim IS, Cheah MH, et al. Structural dynamics in the water and proton channels of photosystem II during the S<sub>2</sub> to S<sub>3</sub> transition. *Nat Commun*. 2021;12(1):6531. <https://doi.org/10.1038/s41467-021-26781-z>
- Ibrahim M, Fransson T, Chatterjee R, Cheah MH, Hussein R, Lassalle L, Sutherlin KD, Young ID, Fuller FD, Gul S, et al. Untangling the sequence of events during the S<sub>2</sub> → S<sub>3</sub> transition in photosystem II and implications for the water oxidation mechanism. *Proc Natl Acad Sci U S A*. 2020;117(23):12624–12635. <https://doi.org/10.1073/pnas.2000529117>

- Ilík P, Pavlovič A, Kouřil R, Alboresi A, Morosinotto T, Allahverdiyeva Y, Aro EM, Yamamoto H, Shikanai T. Alternative electron transport mediated by flavodiiron proteins is operational in organisms from cyanobacteria up to gymnosperms. *New Phytol.* 2017;214(3):967–972. <https://doi.org/10.1111/nph.14536>
- Ishikita H, Saenger W, Biesiadka J, Loll B, Knapp EW. How photosynthetic reaction centers control oxidation power in chlorophyll pairs P680, P700, and P870. *Proc Natl Acad Sci U S A.* 2006;103(26):9855–9860. <https://doi.org/10.1073/pnas.0601446103>
- Jackson PJ, Hitchcock A, Brindley AA, Dickman MJ, Hunter CN. Absolute quantification of cellular levels of photosynthesis-related proteins in *Synechocystis* sp. PCC 6803. *Photosynth Res.* 2023;155(3):219–245. <https://doi.org/10.1007/s11120-022-00990-z>
- Jiao JA, Chollet R. Light/dark regulation of maize leaf phosphoenolpyruvate carboxylase by in vivo phosphorylation. *Arch Biochem Biophys.* 1988;261(2):409–417. [https://doi.org/10.1016/0003-9861\(88\)90357-8](https://doi.org/10.1016/0003-9861(88)90357-8)
- Joet T, Cournac L, Horvath EM, Medgyesy P, Peltier G. Increased sensitivity of photosynthesis to antimycin A induced by inactivation of the chloroplast *ndhB* gene. Evidence for a participation of the NADH-dehydrogenase complex to cyclic electron flow around photosystem I. *Plant Physiol.* 2001;125(4):1919–1929. <https://doi.org/10.1104/pp.125.4.1919>
- Jokel M, Johnson X, Peltier G, Aro EM, Allahverdiyeva Y. Hunting the main player enabling *Chlamydomonas reinhardtii* growth under fluctuating light. *Plant J.* 2018;94(5):822–835. <https://doi.org/10.1111/tjpi.13897>
- Joliot P, Selles J, Wollman FA, Vermeglio A. High efficient cyclic electron flow and functional supercomplexes in *Chlamydomonas* cells. *Biochim Biophys Acta Bioenerg.* 2022;1863(8):148909. <https://doi.org/10.1016/j.bbabi.2022.148909>
- Kammerer B, Fischer K, Hilpert B, Scubert S, Gutensohn M, Weber A, Flugge UI. Molecular characterization of a carbon transporter in plastids from heterotrophic tissues: the glucose 6-phosphate/phosphate antiporter. *Plant Cell.* 1998;10(1):105–117. <https://doi.org/10.1105/tpc.10.1.105>
- Kanygin A, Milrad Y, Thummala C, Reifschneider K, Baker P, Marco P, Yacoby I, Redding KE. Rewiring photosynthesis: a photosystem I-hydrogenase chimera that makes H<sub>2</sub> in vivo. *Energy Environ Sci.* 2020;13(9):2903–2914. <https://doi.org/10.1039/C9EE03859K>
- Kaye Y, Huang W, Clowez S, Saeoussi S, Idoine A, Sanz-Luque E, Grossman AR. The mitochondrial alternative oxidase from *Chlamydomonas reinhardtii* enables survival in high light. *J Biol Chem.* 2019;294(4):1380–1395. <https://doi.org/10.1074/jbc.RA118.004667>
- Kern J, Alonso-Mori R, Hellmich J, Rosalie T, Hattne J, Laksmono H, Gloeckner C, Echols N, Sierra RG, Sellberg J, et al. Room temperature femtosecond X-ray diffraction of photosystem II microcrystals. *Proc Natl Acad Sci U S A.* 2012;109(25):9721–9726. <https://doi.org/10.1073/pnas.1204598109>
- Kern J, Alonso-Mori R, Tran R, Hattne J, Gildea RJ, Echols N, Gloeckner C, Hellmich J, Laksmono H, Sierra RG, et al. Simultaneous femtosecond X-ray spectroscopy and diffraction of photosystem II at room temperature. *Science.* 2013;340(6131):491–495. <https://doi.org/10.1126/science.1234273>
- Kern J, Tran R, Alonso-Mori R, Koroidov S, Echols N, Hattne J, Ibrahim M, Gul S, Laksmono H, Sierra RG, et al. Taking snapshots of photosynthetic water oxidation using femtosecond X-ray diffraction and spectroscopy. *Nature Commun.* 2014;5(1):4371. <https://doi.org/10.1038/ncomms5371>
- Keskin O, Tuncbag N, Gursoy A. Predicting protein–protein interactions from the molecular to the proteome level. *Chem Rev.* 2016;116(8):4884–4909. <https://doi.org/10.1021/acs.chemrev.5b00683>
- Kirchhoff H. Diffusion of molecules and macromolecules in thylakoid membranes. *Biochim Biophys Acta.* 2014a;1837(4):495–502. <https://doi.org/10.1016/j.bbabi.2013.11.003>
- Kirchhoff H. Structural changes of the thylakoid membrane network induced by high light stress in plant chloroplasts. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 2014b;369(1640):20130225. <https://doi.org/10.1098/rstb.2013.0225>
- Kirchhoff H, Haase W, Wegner S, Danielsson R, Ackermann R, Albertsson P-A. Low-light-induced formation of semicrystalline photosystem II arrays in higher plant chloroplasts. *Biochem.* 2007;46(39):11169–11176. <https://doi.org/10.1021/bi700748y>
- Kirchhoff H, Haferkamp S, Allen JF, Epstein DBA, Mullineaux CW. Protein diffusion and macromolecular crowding in thylakoid membranes. *Plant Physiol.* 2008;146(4):1571–1578. <https://doi.org/10.1104/pp.107.115170>
- Kirchhoff H, Mukherjee U, Galla HJ. Molecular architecture of the thylakoid membrane: lipid diffusion space for plastoquinone. *Biochem.* 2002;41(15):4872–4882. <https://doi.org/10.1021/bi011650y>
- Kok B, Forbush B, McGloin M. Cooperation of charges in photosynthetic oxygen evolution. A linear four step mechanism. *Photochem Photobiol.* 1970;11(6):457–475. <https://doi.org/10.1111/j.1751-1097.1970.tb06017.x>
- Komenda J, Sobotka R, Nixon PJ. Assembling and maintaining the photosystem II complex in chloroplasts and cyanobacteria. *Curr Opin Plant Biol.* 2012;15(3):245–251. <https://doi.org/10.1016/j.pbi.2012.01.017>
- Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP. Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science.* 2016;354(6314):857–861. <https://doi.org/10.1126/science.aai8878>
- Kumazawa M, Nishide H, Nagao R, Inoue-Kashino N, Shen J-R, Nakano T, Uchiyama I, Kashino Y, Ifuku K. Molecular phylogeny of fucoxanthin-chlorophyll *a/c* proteins from *Chaetoceros gracilis* and *Lhcq/Lhcf* diversity. *Physiol. Plantarum.* 2022;174(1):e13598. <https://doi.org/10.1111/ppl.13598>
- Lavaud J, Rousseau B, van Gorkom HJ, Etienne A-L. Influence of the diadinoxanthin pool size on photoprotection in the marine planktonic diatom *Phaeodactylum tricornutum*. *Plant Physiol.* 2002;129(3):1398–1406. <https://doi.org/10.1104/pp.002014>
- Lemaire C, Wollman FA, Bennoun P. Restoration of phototrophic growth in a mutant of *Chlamydomonas reinhardtii* in which the chloroplast *atpB* gene of the ATP synthase has a deletion: an example of mitochondria-dependent photosynthesis. *Proc Natl Acad Sci U S A.* 1988;85(5):1344–1348. <https://doi.org/10.1073/pnas.85.5.1344>
- Lemonnier P, Lawson T. Calvin cycle and guard cell metabolism impact stomatal function. *Semin Cell Dev Biol.* 2023;155(Pt A):59–70. <https://doi.org/10.1016/j.semcdb.2023.03.001>
- Levitani O, Chen M, Kuang X, Cheong KY, Jiang J, Banal M, Nambiar N, Gorbunov MY, Ludtke SJ, Falkowski PG, et al. Structural and functional analyses of photosystem II in the marine diatom *Phaeodactylum tricornutum*. *Proc Natl Acad Sci U S A.* 2019;116(35):17316–17322. <https://doi.org/10.1073/pnas.1906726116>
- Li M, Ma J, Li X, Sui S-F. In situ cryo-ET structure of phycobilisome-photosystem II supercomplex from red alga. *eLife.* 2021a;10:e69635. <https://doi.org/10.7554/eLife.69635>
- Li M, Svoboda V, Davis G, Kramer D, Kunz H-H, Kirchhoff H. Impact of ion fluxes across thylakoid membranes on photosynthetic electron transport and photoprotection. *Nat Plants.* 2021b;7(7):979–988. <https://doi.org/10.1038/s41477-021-00947-5>
- Lim SL, Flüttsch S, Liu J, Distefano L, Santelia D, Lim BL. Guard cell chloroplasts import cytosolic ATP for starch turnover and

- stomatal opening. *Nat Commun.* 2022;13(1):652. <https://doi.org/10.1038/s41467-022-28263-2>
- Lindahl M, Spetea C, Hundal T, Oppenheim AB, Adam Z, Andersson B. The thylakoid FtsH protease plays a role in the light-induced turnover of the photosystem II D1 protein. *Plant Cell.* 2000;12(3):419–431. <https://doi.org/10.1105/tpc.12.3.419>
- Liu Y, Mauve C, Lamothe-Sibold M, Guérard F, Glab N, Hodges M, Jossier M. Photorespiratory serine hydroxymethyltransferase 1 activity impacts abiotic stress tolerance and stomatal closure. *Plant Cell Environ.* 2019;42(9):2567–2583. <https://doi.org/10.1111/pce.13595>
- López-Calcano PE, Fisk S, Brown KL, Bull SE, South PF, Raines CA. Overexpressing the H-protein of the glycine cleavage system increases biomass yield in glasshouse and field-grown transgenic tobacco plants. *Plant Biotechnol J.* 2019;17(1):141–151. <https://doi.org/10.1111/pbi.12953>
- Lu Y, Li Y, Yang Q, Zhang Z, Chen Y, Zhang S, Peng XX. Suppression of glycolate oxidase causes glyoxylate accumulation that inhibits photosynthesis through deactivating rubisco in rice. *Physiol Plant.* 2014;150(3):463–476. <https://doi.org/10.1111/pp.12104>
- Lubitiz W, Chrysinina M, Cox N. Water oxidation in photosystem II. *Photosynth. Res.* 2019;142(1):105–125. <https://doi.org/10.1007/s11120-019-00648-3>
- Lüer L, Moulisová V, Henry S, Polli D, Brotosudarmo THP, Hoseinkhani S, Brida D, Lanzani G, Cerullo G, Cogdell RJ. Tracking energy transfer between light harvesting complex 2 and 1 in photosynthetic membranes grown under high and low illumination. *Proc Natl Acad Sci U S A.* 2012;109(5):1473–1478. <https://doi.org/10.1073/pnas.1113080109>
- Lyu MA, Gowik U, Kelly S, Covshoff S, Hibberd JM, Sage RF, Ludwig M, Wong GK, Westhoff P, Zhu XG. The coordination of major events in C(4) photosynthesis evolution in the genus *Flaveria*. *Scientific Rep.* 2021;11(1):15618. <https://doi.org/10.1038/s41598-021-93381-8>
- MacGregor-Chatwin C, Nürnberg DJ, Jackson PJ, Vasilev C, Hitchcock A, Ho M-Y, Shen G, Gisriel CJ, Wood William HJ, Mahbub M. Changes in supramolecular organization of cyanobacterial thylakoid membrane complexes in response to far-red light photoacclimation. *Sci Adv.* 2022;8(6). <https://doi.org/10.1126/sciadv.abj4437>
- MacGregor-Chatwin C, Sener M, Barnett SFH, Hitchcock A, Barnhart-Dailey MC, Maghlaoui K, Barber J, Timlin JA, Schulten K, Hunter CN. Lateral segregation of photosystem I in cyanobacterial thylakoids. *Plant Cell.* 2017;29(5):1119–1136. <https://doi.org/10.1105/tpc.17.00071>
- Mahbub M, Hemm L, Yang Y, Kaur R, Carmen H, Engl C, Huokko T, Riediger M, Watanabe S, Liu L-N, et al. mRNA localization, reaction centre biogenesis and thylakoid membrane targeting in cyanobacteria. *Nat Plants.* 2020;6(9):1179–1191. <https://doi.org/10.1038/s41477-020-00764-2>
- Mahbub M, Mullineaux CW. Locations of membrane protein production in a cyanobacterium. *J Bacteriol.* 2023;205(10):e00209–e00223. <https://doi.org/10.1128/jb.00209-23>
- Maier A, Fahnenstich H, von Caemmerer S, Engqvist MKM, Weber APM, Flügge U-I, Maurino VG. Transgenic introduction of a glycolate oxidative cycle into *A. thaliana* chloroplasts leads to growth improvement. *Front Plant Sci.* 2012;3:38. <https://doi.org/10.3389/fpls.2012.00038>
- Maier A, Zell MB, Maurino VG. Malate decarboxylases: evolution and roles of NAD(P)-ME isoforms in species performing C(4) and C(3) photosynthesis. *J Exp Bot.* 2011;62(9):3061–3069. <https://doi.org/10.1093/jxb/err024>
- Makino A, Miyake C, Yokota A. Physiological functions of the water-water cycle (Mehler reaction) and the cyclic electron flow around PSI in rice leaves. *Plant Cell Physiol.* 2002;43(9):1017–1026. <https://doi.org/10.1093/pcp/pcf124>
- Malone LA, Proctor MS, Hitchcock A, Hunter CN, Johnson MP. Cytochrome b6f—orchestrator of photosynthetic electron transfer. *Biochim Biophys Acta.* 2021;1862(5):148380. <https://doi.org/10.1016/j.bbabi.2021.148380>
- Manna P, Hoffmann M, Davies T, Richardson KH, Johnson MP, Schlau-Cohen GS. Energetic driving force for LHCII clustering in plant membranes. *Sci Adv.* 2023;9(51):2023–2004. <https://doi.org/10.1126/sciadv.adj0807>
- Mao R, Zhang H, Bie L, Liu L-N, Gao J. Million-atom molecular dynamics simulations reveal the interfacial interactions and assembly of plant PSII-LHCII supercomplex. *RSC Adv.* 2023;13(10):6699–6712. <https://doi.org/10.1039/D2RA08240C>
- Mazor Y, Nataf D, Toporik H, Nelson N. Crystal structures of virus-like photosystem I complexes from the mesophilic cyanobacterium *Synechocystis* PCC 6803. *eLife.* 2014;3:e01496. <https://doi.org/10.7554/eLife.01496>
- Messant M, Timm S, Fantuzzi A, Weckwerth W, Bauwe H, Rutherford AW, Krieger-Liszak A. Glycolate induces redox tuning of photosystem II in vivo: study of a photorespiration mutant. *Plant Physiol.* 2018;177(3):1277–1285. <https://doi.org/10.1104/pp.18.00341>
- Mettler T, Mühlhaus T, Hemme D, Schöttler M-A, Rupprecht J, Idoine A, Veyel D, Pal SK, Yaneva-Roder L, Winck FV, et al. Systems analysis of the response of photosynthesis, metabolism, and growth to an increase in irradiance in the photosynthetic model organism *Chlamydomonas reinhardtii*. *Plant Cell* 2014;26(6): 2310–2350 <https://doi.org/10.1105/tpc.114.124537>
- Meyers BE, Boyd SP. Signal decomposition using masked proximal operators. *Found Trend Signal Proc.* 2023;17(1):1–78. <https://doi.org/10.1561/2000000122>
- Mirkovic T, Ostroumov EE, Anna JM, van Grondelle R, Govindjee, Scholes GD. Light absorption and energy transfer in the antenna complexes of photosynthetic organisms. *Chem Rev.* 2017;117(2):249–293. <https://doi.org/10.1021/acs.chemrev.6b00002>
- Morales A, Kaiser E. Photosynthetic acclimation to fluctuating irradiance in plants. *Front Plant Sci.* 2020;11:268. <https://doi.org/10.3389/fpls.2020.00268>
- Munekage Y, Hashimoto M, Miyake C, Tomizawa K, Endo T, Tasaka M, Shikanai T. Cyclic electron flow around photosystem I is essential for photosynthesis. *Nature.* 2004;429(6991):579–582. <https://doi.org/10.1038/nature02598>
- Mustárdy L, Buttle K, Steinbach G, Garab G. The three-dimensional network of the thylakoid membranes in plants: quasi-helical model of the granum-stroma assembly. *Plant Cell.* 2008;20(10):2552–2557. <https://doi.org/10.1105/tpc.108.059147>
- Mustila H, Paananen P, Battchikova N, Santana-Sánchez A, Muth-Pawlak D, Hagemann M, Aro EM, Allahverdiyeva Y. The flavodiiron protein Flv3 functions as a homo-oligomer during stress acclimation and is distinct from the Flv1/Flv3 hetero-oligomer specific to the O<sub>2</sub> photoreduction pathway. *Plant Cell Physiol.* 2016;57(7):1468–1483. <https://doi.org/10.1093/pcp/pcw047>
- Nagao R, Kato K, Ifuku K, Suzuki T, Kumazawa M, Uchiyama I, Kashino Y, Dohmae N, Akimoto S, Shen J-R, et al. Structural basis for assembly and function of a diatom photosystem I-light-harvesting supercomplex. *Nat Commun.* 2020;11(1):2481. <https://doi.org/10.1038/s41467-020-16324-3>
- Nagao R, Kitazaki S, Noguchi T. Evaluation of photosynthetic activities in thylakoid membranes by means of Fourier transform infrared spectroscopy. *Biochim Biophys Acta Bioenerg.* 2018;1859(2):129–136. <https://doi.org/10.1016/j.bbabi.2017.11.004>
- Narusaka Y, Narusaka M, Kobayashi H, Satoh K. The herbicide-resistant species of the cyanobacterial D1 protein obtained by

- thorough and random in vitro mutagenesis. *Plant Cell Physiol.* 1998;39(6):620–626. <https://doi.org/10.1093/oxfordjournals.pcp.a029413>
- Narusaka Y, Narusaka M, Satoh K, Kobayashi H. In vitro random mutagenesis of the D1 protein of the photosystem II reaction center confers phototolerance on the cyanobacterium *Synechocystis* sp. PCC 6803\*. *J Biol Chem.* 1999;274(33):23270–23275. <https://doi.org/10.1074/jbc.274.33.23270>
- Nawrocki WJ, Buchert F, Joliot P, Rappaport F, Bailleul B, Wollman F-A. Chlororespiration controls growth under intermittent light. *Plant Physiol.* 2019;179(2):630–639. [10.1104/pp.18.01213](https://doi.org/10.1104/pp.18.01213)
- Nayak L, Panda D, Dash GK, Lal MK, Swain P, Baig MJ, Kumar A. A chloroplast glycolate catabolic pathway bypassing the endogenous photorespiratory cycle enhances photosynthesis, biomass and yield in rice (*Oryza sativa* L). *Plant Sci.* 2022;314:111103. <https://doi.org/10.1016/j.plantsci.2021.111103>
- Nedbal L, Březina V. Complex metabolic oscillations in plants forced by harmonic irradiance. *Biophys J.* 2002;83(4):2180–2189. [https://doi.org/10.1016/S0006-3495\(02\)73978-7](https://doi.org/10.1016/S0006-3495(02)73978-7)
- Nedbal L, Lazár D. Photosynthesis dynamics and regulation sensed in the frequency domain. *Plant Physiol.* 2021;187(2):646–661. <https://doi.org/10.1093/plphys/kiab317>
- Nelson N, Junge W. Structure and energy transfer in photosystems of oxygenic photosynthesis. *Annu Rev Biochem.* 2015;84(1):659–683. <https://doi.org/10.1146/annurev-biochem-092914-041942>
- Nelson N, Yocum CF. Structure and function of photosystems I and II. *Ann Rev Plant Biol.* 2006;57(1):521–565. <https://doi.org/10.1146/annurev.arplant.57.032905.105350>
- Nevo R, Charuvi D, Shimoni E, Schwarz R, Kaplan A, Ohad I, Reich Z. Thylakoid membrane perforations and connectivity enable intracellular traffic in cyanobacteria. *EMBO J.* 2007;26(5):1467–1473. <https://doi.org/10.1038/sj.emboj.7601594>
- Nguyen TB, Lefoulon C, Nguyen TH, Blatt MR, Carroll W. Engineering stomata for enhanced carbon capture and water-use efficiency. *Trends Plant Sci.* 2023;28(11):1290–1309. <https://doi.org/10.1016/j.tplants.2023.06.002>
- Nicol L, Croce R. The PsbS protein and low pH are necessary and sufficient to induce quenching in the light-harvesting complex of plants LHCII. *Sci Rep.* 2021;11(1):7415. <https://doi.org/10.1038/s41598-021-86975-9>
- Nikkanen L, Santana Sánchez A, Ermakova M, Rögner M, Cournac L, Allahverdiyeva Y. Functional redundancy between flavodiiron proteins and NDH-1 in *Synechocystis* sp. PCC 6803. *Plant J.* 2020;103(4):1460–1476. <https://doi.org/10.1111/tpj.14812>
- Nikkanen L, Vakal S, Santana Sánchez A, Hubáček M, Wang Y, Böhm M, Gutekunst K, Salminen TA, Allahverdiyeva Y. Proton motive force dissipation drives flavodiiron proteins to the thylakoid membrane for ferredoxin-powered O<sub>2</sub> photoreduction. *bioRxiv* 541409, <https://doi.org/10.1101/2023.05.19.541409>, 22 May 2023, preprint: not peer reviewed.
- Niu Y, Lazár D, Holzwarth AR, Kramer DM, Matsubara S, Fiorani F, Poorter H, Schrey SD, Nedbal L. Plants cope with fluctuating light by frequency-dependent nonphotochemical quenching and cyclic electron transport. *New Phytol.* 2023;239(5):1869–1886. <https://doi.org/10.1111/nph.19083>
- Niyogi KK, Truong TB. Evolution of flexible non-photochemical quenching mechanisms that regulate light harvesting in oxygenic photosynthesis. *Curr Opin Plant Biol.* 2013;16(3):307–314. <https://doi.org/10.1016/j.pbi.2013.03.011>
- Nunes-Nesi A, Carrari F, Gibon Y, Sulpice R, Lytovchenko A, Fisahn J, Graham J, Ratcliffe RG, Sweetlove LJ, Fernie AR. Deficiency of mitochondrial fumarate activity in tomato plants impairs photosynthesis via an effect on stomatal function. *Plant J.* 2007;50(6):1093–1106. <https://doi.org/10.1111/j.1365-3113.2007.03115.x>
- Ojakian GK, Satir P. Particle movements in chloroplast membranes: quantitative measurements of membrane fluidity by the freeze-fracture technique. *Proc Natl Acad Sci U S A.* 1974;71(5):2052–2056. <https://doi.org/10.1073/pnas.71.5.2052>
- Oliver T, Kim TD, Trinugroho JP, Córdón-Preciado V, Wijayatilake N, Bhatia A, Rutherford AW, Cardona T. The evolution and evolvability of photosystem II. *Annu Rev Plant Biol.* 2023;74(1):225–257. <https://doi.org/10.1146/annurev-arplant-070522-062509>
- Onoa B, Fukuda S, Iwai M, Bustamante C, Niyogi KK. Atomic force microscopy visualizes mobility of photosynthetic proteins in grana thylakoid membranes. *Biophys J.* 2020;118(8):1876–1886. <https://doi.org/10.1016/j.bpj.2020.02.029>
- Opatíková M, Semchonok DA, Kopečný D, Ilík P, Pospíšil P, Ilíková I, Roudnický P, Zeljković SC, Tarkowski P, Kyrilís FL, et al. Cryo-EM structure of a plant photosystem II supercomplex with light-harvesting protein Lhcb8 and  $\alpha$ -tocopherol. *Nat Plants.* 2023;9(8):1359–1369. <https://doi.org/10.1038/s41477-023-01483-0>
- Papagiannakis E, van Stokkum IHM, Fey H, Büchel C, van Grondelle R. Spectroscopic characterization of the excitation energy transfer in the fucoxanthin-chlorophyll protein of diatoms. *Photosynth Res.* 2005;86(1–2):241–250. <https://doi.org/10.1007/s11220-005-1003-8>
- Park S, Steen CJ, Fischer AL, Fleming GR. Snapshot transient absorption spectroscopy: toward in vivo investigations of nonphotochemical quenching mechanisms. *Photosyn Res.* 2019;141(3):367–376. <https://doi.org/10.1007/s11220-019-00640-x>
- Pearcy RW. Sunflecks and photosynthesis in plant canopies. *Annu Rev Plant Physiol Plant Mol Biol.* 1990;41(1):421–453. <https://doi.org/10.1146/annurev.pp.41.060190.002225>
- Peltier G, Aro EM, Shikanai T. NDH-1 and NDH-2 plastoquinone reductases in oxygenic photosynthesis. *Annu Rev Plant Biol.* 2016;67(1):55–80. <https://doi.org/10.1146/annurev-arplant-043014-114752>
- Peltier G, Stoffel C, Findinier J, Madireddi SK, Dao O, Epting V, Morin A, Grossman A, Li-Beisson Y, Burlacot A. Green algal CO<sub>2</sub> capture is powered by alternative electron pathways of photosynthesis. *bioRxiv* 552514, <https://doi.org/10.1101/2023.08.08.552514>, 22 November 2023, preprint: not peer reviewed.
- Peltier G, Tolleter D, Billon E, Cournac L. Auxiliary electron transport pathways in chloroplasts of microalgae. *Photosynth Res.* 2010;106(1–2):19–31. <https://doi.org/10.1007/s11220-010-9575-3>
- Peng J, Lu J, Hoh D, Dina AS, Shang X, Kramer DM, Chen J. Identifying emerging phenomenon in long temporal phenotyping experiments. *Bioinformatics.* 2020;36(2):568–577. <https://doi.org/10.1093/bioinformatics/btz559>
- Perera-Castro AV, Flexas J. Recent advances in understanding and improving photosynthesis. *Fac Rev.* 2020;9:5. <https://doi.org/10.12703/b/9-5>
- Perez-Boerema A, Engel BD, Wietrzynski W. Evolution of thylakoid structural diversity. *Annu Rev Cell Dev Biol.* <https://doi.org/10.1146/annurev-cellbio-120823-022747>, 2024, preprint: not peer reviewed.
- Perin G, Bellan A, Bernardi A, Bezzo F, Morosinotto T. The potential of quantitative models to improve microalgae photosynthetic efficiency. *Physiol Plantarum.* 2019;166(1):380–391. <https://doi.org/10.1111/ppl.12915>
- Perin G, Bellan A, Michelberger T, Lyska D, Wakao S, Niyogi KK, Morosinotto T. Modulation of xanthophyll cycle impacts biomass productivity in the marine microalga *nannochloropsis*. *Proc. Natl.*

- Acad. Sci. U S A. 2023;120(25):e2214119120. [10.1073/pnas.2214119120](https://doi.org/10.1073/pnas.2214119120)
- Petrou K, Belgio E, Ruban AV. Ph sensitivity of chlorophyll fluorescence quenching is determined by the detergent/protein ratio and the state of LHCII aggregation. *Biochim Biophys Acta*. 2014;1837(9):1533–1539. <https://doi.org/10.1016/j.bbabi.2013.11.018>
- Pi X, Zhao S, Wang W, Liu D, Xu C, Han G, Kuang T, Sui S-F, Shen J-R. The pigment-protein network of a diatom photosystem II-light-harvesting antenna supercomplex. *Science*. 2019;365(6452):eaax4406. <https://doi.org/10.1126/science.aax4406>
- Prabhakar V, Lottgert T, Geimer S, Dormann P, Kruger S, Vijayakumar V, Schreiber L, Gobel C, Feussner K, Freussner I, et al. Phosphoenolpyruvate provision to plastids is essential for gametophyte and sporophyte development in *Arabidopsis thaliana*. *Plant Cell*. 2010;22(8):2594–2617. <https://doi.org/10.1105/tpc.109.073171>
- Rast A, Schaffer M, Albert S, Wan W, Pfeffer S, Beck F, Plitzko JM, Nickelsen J, Engel BD. Biogenic regions of cyanobacterial thylakoids form contact sites with the plasma membrane. *Nat Plants*. 2019;5(4):436–446. <https://doi.org/10.1038/s41477-019-0399-7>
- Rea G, Lambrea M, Politicelli F, Bertalan I, Antonacci A, Pastorelli S, Damasso M, Johanningmeier U, Giardi MT. Directed evolution and in silico analysis of reaction centre proteins reveal molecular signatures of photosynthesis adaptation to radiation pressure. *PLoS One*. 2011;6(1):e16216. <https://doi.org/10.1371/journal.pone.0016216>
- Roach T. LHCSR3-Type NPQ prevents photoinhibition and slowed growth under fluctuating light in *Chlamydomonas reinhardtii*. *Plants*. 2020;9(11):1604. <https://doi.org/10.3390/plants9111604>
- Rödiger A, Boekema E, Büchel C. The structure of FCPb, a light-harvesting complex in the diatom *Cyclotella meneghiniana*. *Photosynth Res*. 2018;135(1–3):203–211. <https://doi.org/10.1007/s11120-016-0328-9>
- Rogov AG, Sukhanova EI, Uralskaya LA, Aliverdieva DA, Zvyagil'skaya RA. Alternative oxidase: distribution, induction, properties, structure, regulation, and functions. *Biochemistry (Moscow)*. 2014;79(13):1615–1634. <https://doi.org/10.1134/S0006297914130112>
- Rosnow JJ, Edwards GE, Roalson EH. Positive selection of Kranz and non-Kranz C4 phosphoenolpyruvate carboxylase amino acids in Suaedoideae (Chenopodiaceae). *J Exp Bot*. 2014;65(13):3595–3607. <https://doi.org/10.1093/jxb/eru053>
- Ruban AV, Johnson MP. Visualizing the dynamic structure of the plant photosynthetic membrane. *Nat Plants*. 2015;1(11):15161. <https://doi.org/10.1038/nplants.2015.161>
- Ruban AV, Lavaud J, Rousseau B, Guglielmi G, Horton P, Etienne A-L. The super-excess energy dissipation in diatom algae: comparative analysis with higher plants. *Photosynth Res*. 2004;82(2):165–175. <https://doi.org/10.1007/s11120-004-1456-1>
- Sage RF, Zhu XG. Exploiting the engine of C(4) photosynthesis. *J Exp Bot*. 2011;62(9):2989–3000. <https://doi.org/10.1093/jxb/err179>
- Saito S, Uozumi N. Guard cell membrane anion transport systems and their regulatory components: an elaborate mechanism controlling stress-induced stomatal closure. *Plants*. 2019;8(1):9. <https://doi.org/10.3390/plants8010009>
- Santana-Sánchez A, Nikkanen L, Werner E, Tóth G, Ermakova M, Kosourov S, Walter J, He M, Aro EM, Allahverdiyeva Y. Flv3A facilitates O<sub>2</sub> photoreduction and affects H<sub>2</sub> photoproduction independently of Flv1A in diazotrophic *Anabaena* filaments. *New Phytol*. 2023;237(1):126–139. <https://doi.org/10.1111/nph.18506>
- Santana-Sánchez A, Solymosi D, Mustila H, Bersanini L, Aro EM, Allahverdiyeva Y. Flavodiiron proteins 1-to-4 function in versatile combinations in O<sub>2</sub> photoreduction in cyanobacteria. *Elife*. 2019;8:e45766. <https://doi.org/10.7554/eLife.45766>
- Sarcina M, Bouzovitis N, Mullineaux CW. Mobilization of photosystem II induced by intense red light in the cyanobacterium *Synechococcus* sp PCC7942. *Plant Cell*. 2006;18(2):457–464. <https://doi.org/10.1105/tpc.105.035808>
- Saroussi S, Sanz-Luque E, Kim RG, Grossman AR. Nutrient scavenging and energy management: acclimation responses in nitrogen and sulfur deprived *Chlamydomonas*. *Curr Opin Plant Biol*. 2017;39:114–122. <https://doi.org/10.1016/j.pbi.2017.06.002>
- Saroussi SI, Wittkopp TM, Grossman AR. The type II NADPH dehydrogenase facilitates cyclic electron flow, energy-dependent quenching, and chlororespiratory metabolism during acclimation of *Chlamydomonas reinhardtii* to nitrogen deprivation. *Plant Physiol*. 2016;170(4):1975–1988. <https://doi.org/10.1104/pp.15.02014>
- Schlau-Cohen GS, Yang H-Y, Krüger TPJ, Xu P, Gwizdala M, van Grondelle R, Croce R, Moerner WE. Single-molecule identification of quenched and unquenched states of LHCII. *J Phys Chem Lett*. 2015;6(5):860–867. <https://doi.org/10.1021/acs.jpcclett.5b00034>
- Schneider AR, Geissler PR. Coexistence of fluid and crystalline phases of proteins in photosynthetic membranes. *Biophys J*. 2013;105(5):1161–1170. <https://doi.org/10.1016/j.bpj.2013.06.052>
- Schottkowski M, Peters M, Zhan Y, Rifai O, Zhang Y, Zerges W. Biogenic membranes of the chloroplast in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci U S A*. 2012;109(47):19286–19291. <https://doi.org/10.1073/pnas.1209860109>
- Schubert H, Sagert S, Forster RM. Evaluation of the different levels of variability in the underwater light field of a shallow estuary. *Helgoland Marine Research*. 2001;55(1):12–22. <https://doi.org/10.1007/s101520000064>
- Sejima T, Takagi D, Fukayama H, Makino A, Miyake C. Repetitive short-pulse light mainly inactivates photosystem I in sunflower leaves. *Plant Cell Physiol*. 2014;55(6):1184–1193. <https://doi.org/10.1093/pcp/pcu061>
- Sello S, Meneghesso A, Alboresi A, Baldan B, Morosinotto T. Plant biodiversity and regulation of photosynthesis in the natural environment. *Planta*. 2019;249(4):1217–1228. <https://doi.org/10.1007/s00425-018-03077-z>
- Sétif P, Shimakawa G, Krieger-Liszakay A, Miyake C. Identification of the electron donor to flavodiiron proteins in *Synechocystis* sp. PCC 6803 by in vivo spectroscopy. *Biochim Biophys Acta Bioenerg*. 2020;1861(10):148256. <https://doi.org/10.1016/j.bbabi.2020.148256>
- Sharkey TD. The discovery of rubisco. *J Exp Bot*. 2022;74(2):510–519. <https://doi.org/10.1093/jxb/erac254>
- Sharon I, Alperovitch A, Rohwer F, Haynes M, Glaser F, Atamna-Ismael N, Pinter RY, Partensky F, Koonin EV, Wolf YI, et al. Photosystem I gene cassettes are present in marine virus genomes. *Nature*. 2009;461(7261):258–262. <https://doi.org/10.1038/nature08284>
- Shi T, Bibby TS, Jiang L, Irwin AJ, Falkowski PG. Protein interactions limit the rate of evolution of photosynthetic genes in Cyanobacteria. *Mol Biol Evol*. 2005;22(11):2179–2189. <https://doi.org/10.1093/molbev/msi216>
- Shikanai T, Endo T, Hashimoto T, Yamada Y, Asada K, Yokota A. Directed disruption of the tobacco ndhB gene impairs cyclic electron flow around photosystem I. *Proc Natl Acad Sci U S A*. 1998;95(16):9705–9709. <https://doi.org/10.1073/pnas.95.16.9705>
- Shimakawa G, Miyake C. Changing frequency of fluctuating light reveals the molecular mechanism for P700 oxidation in plant leaves. *Plant Direct*. 2018;2(7):e00073. <https://doi.org/10.1002/pld3.73>
- Shimakawa G, Shaku K, Nishi A, Hayashi R, Yamamoto H, Sakamoto K, Makino A, Miyake C. FLAVODIIRON2 and FLAVODIIRON4

- proteins mediate an oxygen-dependent alternative electron flow in *Synechocystis* sp. PCC 6803 under CO<sub>2</sub>-limited conditions. *Plant Physiol.* 2015;167(2):472–480. <https://doi.org/10.1104/pp.114.249987>
- Shimoni E, Rav-Hon O, Ohad I, Brumfeld V, Reich Z. Three-dimensional organization of higher-plant chloroplast thylakoid membranes revealed by electron tomography. *Plant Cell.* 2005;17(9):2580–2586. <https://doi.org/10.1105/tpc.105.035030>
- Smith WK, Berry ZC. Sunflecks? *Tree Physiol.* 2013;33(3):233–237. <https://doi.org/10.1093/treephys/tpt005>
- Son M, Moya R, Pinnola A, Bassi R, Schlau-Cohen GS. Protein–protein interactions induce pH-dependent and zeaxanthin-independent photoprotection in the plant light-harvesting complex, LHCI. *J Am Chem Soc.* 2021;143(42):17577–17586. <https://doi.org/10.1021/jacs.1c07385>
- South PF, Cavanagh AP, Liu HW, Ort DR. Synthetic glycolate metabolism pathways stimulate crop growth and productivity in the field. *Science.* 2019;363(6422):eaat9077. <https://doi.org/10.1126/science.aat9077>
- Staelin LA. Reversible particle movements associated with unstacking and restacking of chloroplast membranes in vitro. *J Cell Biol.* 1976;71(1):136–158. <https://doi.org/10.1083/jcb.71.1.136>
- Standfuss J, van Scheltinga ACT, Lamborghini M, Kühlbrandt W. Mechanisms of photoprotection and nonphotochemical quenching in pea light-harvesting complex at 2.5 Å resolution. *EMBO J.* 2005;24(5):919–928. <https://doi.org/10.1038/sj.emboj.7600585>
- Steen CJ, Burlacot A, Short AH, Niyogi KK, Fleming GR. Interplay between LHCSR proteins and state transitions governs the NPQ response in intact cells of *Chlamydomonas* during light fluctuations. *Plant Cell Environ.* 2022;45(8):2428–2445. <https://doi.org/10.1111/pce.14372>
- Stirbet A, Lázár D, Guo Y, Govindjee G. Photosynthesis: basics, history and modelling. *Ann Bot.* 2020;126(4):511–537. <https://doi.org/10.1093/aob/mcz171>
- Storti M, Alborese A, Gerotto C, Aro E-M, Finazzi G, Morosinotto T. Role of cyclic and pseudo-cyclic electron transport in response to dynamic light changes in *Physcomitrella patens*. *Plant Cell Environ.* 2019;42(5):1590–1602. <https://doi.org/10.1111/pce.13493>
- Streatfield SJ, Weber A, Kinsman EA, Hausler RE, Li J, Post-Beittenmiller D, Kaiser WM, Pyke KA, Flugge UI, Chory J. The phosphoenolpyruvate/phosphate translocator is required for phenolic metabolism, palisade cell development, and plastid-dependent nuclear gene expression. *Plant Cell.* 1999;11(9):1609–1622. <https://doi.org/10.1105/tpc.11.9.1609>
- Suga M, Akita F, Hirata K, Ueno G, Murakami H, Nakajima Y, Shimizu T, Yamashita K, Yamamoto M, Ago H, et al. Native structure of photosystem II at 1.95 Å resolution viewed by femtosecond X-ray pulses. *Nature.* 2015;517(7532):99–103. <https://doi.org/10.1038/nature13991>
- Sznee K, Dekker JP, Dame RT, van Roon H, Wuite GJL, Frese RN. Jumping mode atomic force microscopy on grana membranes from spinach. *J Biol Chem.* 2011;286(45):39164–39171. <https://doi.org/10.1074/jbc.M111.284844>
- Taddei L, Stella GR, Rogato A, Bailleul B, Fortunato AE, Annunziata R, Sanges R, Thaler M, Lepetit B, Lavaud J, et al. Multisignal control of expression of the LHGX protein family in the marine diatom *Phaeodactylum tricornutum*. *J Exp Bot.* 2016;67(13):3939–3951. <https://doi.org/10.1093/jxb/erw198>
- Tikkanen M, Grieco M, Kangasjärvi S, Aro E-M. Thylakoid protein phosphorylation in higher plant chloroplasts optimizes electron transfer under fluctuating light. *Plant Physiol.* 2010;152(2):723–735. <https://doi.org/10.1104/pp.109.150250>
- Timm S, Mielewicz M, Florian A, Frankenbach S, Dreissen A, Hocken N, Fernie AR, Walter A, Bauwe H. High-to-low CO<sub>2</sub> acclimation reveals plasticity of the photorespiratory pathway and indicates regulatory links to cellular metabolism of Arabidopsis. *PLoS One.* 2012;7(8):e42809. <https://doi.org/10.1371/journal.pone.0042809>
- Timm S, Woitschach F, Heise C, Hagemann M, Bauwe H. Faster removal of 2-phosphoglycolate through photorespiration improves abiotic stress tolerance of Arabidopsis. *Plants.* 2019;8(12):563. <https://doi.org/10.3390/plants8120563>
- Tiwari V, Matutes YA, Gardiner AT, Jansen TLC, Cogdell RJ, Ogilvie JP. Spatially-resolved fluorescence-detected two-dimensional electronic spectroscopy probes varying excitonic structure in photosynthetic bacteria. *Nat Commun.* 2018;9(1):4219. <https://doi.org/10.1038/s41467-018-06619-x>
- Tronconi MA, Hüdig M, Schranz ME, Maurino VG. Independent recruitment of duplicated β-subunit-coding NAD-ME genes aided the evolution of C4 photosynthesis in Cleomaceae. *Front Plant Sci.* 2020;11:572080. <https://doi.org/10.3389/fpls.2020.572080>
- Tsuchida Y, Furumoto T, Izumida A, Hata S, Izui K. Phosphoenolpyruvate carboxylase kinase involved in C(4) photosynthesis in *Flaveria trinervia*: cDNA cloning and characterization. *FEBS Lett.* 2001;507(3):318–322. [https://doi.org/10.1016/S0014-5793\(01\)02994-5](https://doi.org/10.1016/S0014-5793(01)02994-5)
- Umena Y, Kawakami K, Shen J-R, Kamiya N. Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature.* 2011;473(7345):55–60. <https://doi.org/10.1038/nature09913>
- van Grondelle R, Novoderezhkin VI. Energy transfer in photosynthesis: experimental insights and quantitative models. *Phys Chem Chem Phys.* 2006;8(7):793–807. <https://doi.org/10.1039/B514032C>
- Vicente JB, Carrondo MA, Teixeira M, Frazão C. Structural studies on flavodiiron proteins. *Methods Enzymol.* 2008;437:3–19. [https://doi.org/10.1016/S0076-6879\(07\)37001-8](https://doi.org/10.1016/S0076-6879(07)37001-8)
- von Bismarck T, Korkmaz K, Ruß J, Skurk K, Kaiser E, Correa Galvis V, Cruz JA, Strand DD, Köhl K, Eirich J, et al. Light acclimation interacts with thylakoid ion transport to govern the dynamics of photosynthesis in Arabidopsis. *New Phytol.* 2023;237(1):160–176. <https://doi.org/10.1111/nph.18534>
- Walters RG, Ibrahim DG, Horton P, Kruger NJ. A mutant of Arabidopsis lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch breakdown in the light. *Plant Physiol.* 2004;135(2):891–906. <https://doi.org/10.1104/pp.104.040469>
- Wang D, Fiebig OC, Harris D, Toporik H, Ji Y, Chuang C, Nairat M, Tong AL, Ogren JI, Hart SM, et al. Elucidating interprotein energy transfer dynamics within the antenna network from purple bacteria. *Proc Natl Acad Sci U S A.* 2023a;120(28):e2220477120. <https://doi.org/10.1073/pnas.2220477120>
- Wang P, Frank A, Appel J, Boehm M, Strabel N, Nowaczyk MM, Schuhmann W, Conzuelo F, Gutekunst K. In vivo assembly of photosystem I-hydrogenase chimera for in vitro photoH<sub>2</sub> production. *Adv Energy Mater.* 2023b;13(14):2203232. <https://doi.org/10.1002/aenm.202203232>
- Wang W, Yu L-J, Xu C, Tomizaki T, Zhao S, Umena Y, Chen X, Qin X, Xin Y, Suga M, et al. Structural basis for blue-green light harvesting and energy dissipation in diatoms. *Science.* 2019;363(6427):eaav0365. <https://doi.org/10.1126/science.aav0365>
- Wang W, Zhao S, Shen L, Li X, Tao Q, Xu C, Zhou C, Yang Y, Sang M, Han G, et al. Structural insights into photosystem II supercomplex and trimeric FCP antennae of a centric diatom *Cyclotella meneghiniana*. *Nat Commun.* 2023c;14(1):8164. <https://doi.org/10.1038/s41467-023-30949-2>
- Wietrzynski W, Schaffer M, Tegunov D, Albert S, Kanazawa A, Plietzko JM, Baumeister W, Engel BD. Charting the native architecture of *Chlamydomonas* thylakoid membranes with single-molecule precision. *Elife.* 2020;9:e53740. <https://doi.org/10.7554/eLife.53740>

- Wood WHJ, MacGregor-Chatwin C, Barnett SFH, Mayneord GE, Huang X, Hobbs JK, Hunter CN, Johnson MP. Dynamic thylakoid stacking regulates the balance between linear and cyclic photosynthetic electron transfer. *Nat Plants*. 2018;4(2):116–127. <https://doi.org/10.1038/s41477-017-0092-7>
- Wydrzynski T, Satoh S. *Photosystem II: the light-driven water:plastoquinone oxidoreductase*. Dordrecht (The Netherlands): Springer; 2005.
- Xu C, Pi X, Huang Y, Han G, Chen X, Qin X, Huang G, Zhao S, Yang Y, Kuang T, et al. Structural basis for energy transfer in a huge diatom PSI-FCPI supercomplex. *Nat Commun*. 2020;11(1):5081. <https://doi.org/10.1038/s41467-020-18867-x>
- Yachandra VK, Sauer K, Klein MP. Manganese cluster in photosynthesis: where plants oxidize water to dioxygen. *Chem Rev*. 1996;96(7):2927–2950. <https://doi.org/10.1021/cr950052k>
- Yamamoto H, Peng L, Fukao Y, Shikanai T. An Src homology 3 domain-like fold protein forms a ferredoxin binding site for the chloroplast NADH dehydrogenase-like complex in Arabidopsis. *Plant Cell*. 2011;23(4):1480–1493. <https://doi.org/10.1105/tpc.110.080291>
- Yamamoto H, Shikanai T. PGR5-Dependent cyclic electron flow protects photosystem I under fluctuating light at donor and acceptor sides. *Plant Physiol*. 2018;179(2):588–600. <https://doi.org/10.1104/pp.18.01343>
- Yamamoto H, Takahashi S, Badger MR, Shikanai T. Artificial remodelling of alternative electron flow by flavodiiron proteins in Arabidopsis. *Nat Plants*. 2016;2(3):16012. <https://doi.org/10.1038/nplants.2016.12>
- Yano J, Yachandra VK. Mn4Ca cluster in photosynthesis: where and how water is oxidized to dioxygen. *Chem Rev*. 2014;114(8):4175–4205. <https://doi.org/10.1021/cr4004874>
- Yokochi Y, Yoshida K, Hahn F, Miyagi A, Wakabayashi K-i, Kawai-Yamada M, Weber APM, Hisabori T. Redox regulation of NADP-malate dehydrogenase is vital for land plants under fluctuating light environment. *Proc Natl Acad Sci U S A*. 2021;118(6):e2016903118. <https://doi.org/10.1073/pnas.2016903118>
- You X, Zhang X, Cheng J, Xiao Y, Ma J, Sun S, Zhang X, Wang HW, Sui SF. In situ structure of the red algal phycobilisome–PSII–PSI–LHC megacomplex. *Nature*. 2023;616(7955):199–206. <https://doi.org/10.1038/s41586-023-05831-0>
- Zabret J, Bohn S, Schuller SK, Arnolds O, Möller M, Meier-Credo J, Liauw P, Chan A, Tajkhorshid E, Langer JD, et al. Structural insights into photosystem II assembly. *Nat Plants*. 2021;7(4):524–538. <https://doi.org/10.1038/s41477-021-00895-0>
- Zelitch I, Walker DA. The role of glycolic acid metabolism in opening of leaf stomata. *Plant Physiol*. 1964;39(5):856–862. <https://doi.org/10.1104/pp.39.5.856>
- Zeno G, Gomez RL, Caferra R, Stuttmann J, Dall'Osto, L, Bassi R. Thylakoid grana stacking revealed by multiplex genome editing of LHCII encoding genes. bioRxiv 474624, <https://doi.org/10.1101/2021.12.31.474624>, 1 January 2022, preprint: not peer reviewed.