




ASO Visual Abstract: Postoperative Complications After Neoadjuvant Chemotherapy Compared with Upfront Surgery in Gastric Adenocarcinoma—A Population-Based, Nationwide Study in Finland

Emilia Putila, BM¹ , Olli Helminen, MD, PhD¹, Mika Helmiö, MD, PhD², Heikki Huhta, MD, PhD¹, Aapo Jalkanen, MD³, Raija Kallio, MD, PhD⁴, Vesa Koivukangas, MD, PhD¹, Arto Kokkola, MD, PhD³, Simo Laine, MD, PhD², Elina Lietzen, MD, PhD², Johanna Louhimo, MD, PhD³, Sanna Meriläinen, MD, PhD¹, Vesa-Matti Pohjanen, MD, PhD⁵, Tuomo Rantanen, MD, PhD⁶, Anna Junttila, MD, PhD², Ari Ristimäki, MD, PhD^{7,8}, Jari V. Räsänen, MD, PhD⁹, Juha Saarnio, MD, PhD¹, Eero Sihvo, MD, PhD¹⁰, Vesa Toikkanen, MD, PhD¹¹, Tuula Tyrväinen, MD, PhD¹², Antti Valtola, MD⁶, and Joonas H. Kauppila, MD, PhD^{1,13} on behalf of the FINEGO Group

¹Surgery Research Unit, Department of Surgery, Medical Research Center Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland; ²Division of Digestive Surgery and Urology, Turku University Hospital, Turku, Finland; ³Department of Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland; ⁴Department of Oncology and Radiotherapy, Oulu University Hospital, Oulu, Finland; ⁵Cancer and Translational Medicine Research Unit, Medical Research Center Oulu, University of Oulu and Oulu University Hospital, Oulu, Finland; ⁶Department of Surgery, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland; ⁷Department of Pathology, HUSLAB, HUS Diagnostic Center, Helsinki University Hospital and University of Helsinki, Helsinki, Finland; ⁸Applied Tumor Genomics Research Program, Research Programs Unit, Faculty of Medicine, University of Helsinki, Helsinki, Finland; ⁹Department of General Thoracic and Oesophageal Surgery, Heart and Lung Centre, University of Helsinki and Helsinki University Hospital, Helsinki, Finland; ¹⁰Department of Surgery, Central Finland Central Hospital, Jyväskylä, Finland; ¹¹Department of Cardiothoracic Surgery, Heart Center, Tampere University Hospital and University of Tampere, Tampere, Finland; ¹²Department of Gastroenterology and Alimentary Tract Surgery, Tampere University Hospital, Tampere, Finland; ¹³Department of Molecular Medicine and Surgery, Karolinska Institutet and Karolinska University Stockholm, Stockholm, Sweden

This study (<https://doi.org/10.1245/s10434-023-14813-5>) compares neoadjuvant chemotherapy and upfront surgery after gastrectomy for gastric adenocarcinoma in a population-based setting ($n = 1769$). Neoadjuvant chemotherapy did not increase

major postoperative complications (odds ratio 1.12, 95% confidence interval 0.81–1.56). Furthermore, it did not increase reoperations or short term mortality compared to upfront surgery.



Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.