

Supplementary table 3. Clinical parameters of each subject group: control, generalized periodontitis (GP), and type 2 diabetes mellitus patients with GP (T2DM/GP). Data are presented as mean \pm standard deviation (and min-max values for the probing pocket depth).

	CONTROL (n=11)	GP (n=7)	T2DM/GP (n=11)	<i>p</i> -value		
				Control vs. GP	Control vs. T2DM/GP	GP vs. T2DM/GP
Age (years)	27.1 \pm 11.9	51.8 \pm 6.4	52.6 \pm 9.9	<0.001	<0.001	1.000
Probing pocket depth (in mm)	1.9 \pm 0.4 (1.43-2.78)	4.9 \pm 0.7 (3.69-5.83)	4.8 \pm 0.7 (3.86-5.77)	<0.001	<0.001	1.000
Clinical attachment level (in mm)	1.9 \pm 0.4	6.2 \pm 1.1	5.9 \pm 1.2	<0.001	<0.001	1.000
Gingival index	0.5 \pm 0.2	1.0 \pm 0.3	0.9 \pm 0.2	<0.001	<0.001	1.000
Plaque index	0.5 \pm 0.2	0.9 \pm 0.3	0.8 \pm 0.2	0.003	0.001	0.924
Bleeding time index	0.6 \pm 0.7	2.8 \pm 0.4	2.6 \pm 0.5	<0.001	<0.001	1.000
Number of teeth	27.2 \pm 1.4	21.7 \pm 5.3	21.0 \pm 5.1	0.005	0.033	1.000